

- **Food Preservation with antimicrobial preservatives:**

The use of chemicals to prevent or delay the spoilage of foods derives in part from the fact that such compounds are used in the treatment of diseases of humans, animals, and plants. Many chemical compounds, **either present naturally, formed during processing, or legally added as ingredients**, can kill microorganisms or control their growth in foods. They are, as a group, designated as **preservatives**. Some of the **naturally** occurring preservatives can be present in sufficient amounts in foods to produce antimicrobial action, such as lysozyme in egg white and organic acids in citrus fruits. Some of the antimicrobials can be formed in enough quantities **during food processing** to control undesirable microbial growth, such as lactic acid in yogurt fermentation. Among the many **food additives**, some are specifically used to preserve foods against microorganisms (such as NO<sub>2</sub> in cured meat to control spore germination, especially of *Cl. botulinum*) whereas others are added to improve the functional properties of a food (such as butylhydroxyanisol, BHA, used as an antioxidant, although it has antimicrobial properties).

**\*Ideal Antimicrobial preservatives**

Several factors need to be considered in evaluating the suitability of an antimicrobial agent as a food preservative, based on:

- 1- Antimicrobial properties: A compound with a broader antimicrobial spectrum is more suitable for application in foods so that it is effective against many types of microorganisms important in foods ( molds, yeasts, bacteria, and viruses), as compared with one that has a narrow spectrum. Also, a compound effective not only against vegetative cells but also against spores is preferred. Finally, it should not allow development of resistant strains.
- 2- Suitability for application in a food: A compound should not only have the desired antimicrobial property but also not affect the normal quality of a food (texture, flavor, or color). It should not interact with food components and become inactive. It should have a high antimicrobial property at the pH,

$a_w$ , Eh, and storage temperature of the food. It should be stable during the storage of the food. Finally, it should be economical and readily available.

- 3- Ability to meet regulatory requirements: The regulatory requirements include the effectiveness of an antimicrobial agent in a food system. It should be effective in small concentrations and should not conceal poor quality and spoilage of a food. Most importantly, it should be non toxic and safe for human consumption, generally recognized as safe (GRAS). Finally it should be listed in the label, indicating its purpose in the food.

### **Antimicrobial preservatives added to foods:**

#### **1- Acids**

##### **A- Benzoic acid and parabens**

Benzoic acid and its sodium salt , The antimicrobial activity of benzoate is related to pH, the greatest activity being at low pH values. Benzoic acid/benzoates are incorporated into margarine, pickles, apple cider, soft drinks, tomato catsup, salad dressings. They are inhibit Yeasts and molds.

**Parabens (Esters of *p*-Hydroxy benzoic Acid):** Parabens are used as methyl, ethyl, butyl, or propyl parabens. They are broad spectrum antimicrobial agents, they are effective at high pH and against bacteria, yeasts, and molds. They are used in fruit fillings, jams and jellies, pickles, confectioneries, salad dressings and mustards.

##### **B-Acetic Acid**

Acetic acid is used usually as vinegar (5 to 10% acetic acid) or as salts of sodium and calcium in pickles, salad dressings, and sauces. It is more effective against bacteria than yeasts and molds.

##### **C-Propionic Acid**

Propionic acid is used as salts of calcium and sodium in bread, bakery products, cheeses, jam and jellies, and tomato puree. It is effective against molds and bacteria but almost ineffective against yeasts.

### **D-Lactic Acid**

Lactic acid is used as acid or sodium and potassium salts in carbonated drinks, salad dressings, pickled vegetables, low-heat-processed meat products, and sauces. It is less effective than acetic, propionic, benzoic, or sorbic acids, but more effective than citric acid. It is more effective against bacteria but quite ineffective against yeasts and molds.

### **E-Citric Acid**

Citric acid is used at 1% (or more) in nonalcoholic drinks, jams and jellies, baking products, cheeses, canned vegetables, and sauces. It is less effective than lactic acid against bacteria as well as yeasts and molds.


### **F-Sorbic Acid**

It is an unsaturated acid and used either as acid or as salts of sodium, potassium, or calcium. It is used in nonalcoholic drinks, some alcoholic drinks, processed fruits and vegetables, dairy desserts, confectioneries, mayonnaise, salad dressings and mustards. It is more effective against molds and yeasts than against bacteria. Among bacteria, catalase-negative (e.g., lactic acid bacteria) are more resistant than catalase-positive species (e.g., aerobes, *S. aureus*, and *Bacillus* spp.). Also, aerobic bacteria are more sensitive to it than anaerobic bacteria.

### **2-Nitrite and Nitrate**

Curing agents that contain nitrite, and together with NaCl, sugar, spices, ascorbate, and erythorbate, are permitted for use in heat-processed meat, poultry, and fish products to control growth and toxin production by *Clostridium botulinum*. The antibotulinal effect consists of inhibition of vegetative cell growth and the prevention of germination and growth of spores that survive heat processing or smoking during postprocessing storage.

Nitrate and nitrite are also used in several European countries in some cheeses to prevent gas blowing by *Cl. butyricum* and *Cl. tyrobutyricum*. In addition to clostridial species, nitrite is inhibitory, to some extent, to *Staphylococcus aureus*, *Escherichia*, *Pseudomonas*, and *Enterobacter* spp. It also forms desirable product color and enhances flavor in cured meat products. Nitric oxide (NO), the important product from the standpoint of color fixation in cured meats. Nitric oxide reacts with myoglobin under reducing conditions to produce the desirable red pigment nitrosomyoglobin.



When nitrite reacts with secondary amines, nitrosamines are formed, and many are known to be carcinogenic. Nitrosamines have been found in cured meat and fish products at low levels. Because of this, there is a trend to reduce NO<sub>2</sub> or to use other preservatives to control *Cl.botulinum* in low-heat-processed meat products.

### 3-Sulfur Dioxide and Sulfites

Sulfur dioxide, sodium sulfite (NaSO<sub>3</sub>), sodium bisulfite (NaHSO<sub>3</sub>), and sodium metabisulfite (Na<sub>2</sub>S<sub>2</sub>O<sub>5</sub>) are used to control microorganisms (and insects) in soft fruits, fruit juices, lemon juices, beverages, wines, sausages, pickles, and fresh shrimp. They are more effective against molds and yeasts than bacteria; among bacteria, the aerobic Gram-negative rods are the most susceptible. The antimicrobial action is produced by the undissociated sulfurous acid that rapidly enters the cell and reacts with the thiol groups in structural proteins, enzymes, and cofactors, as well as with other cellular components. Sulfur dioxide and sulfites are also used as antioxidants in fresh and dried fruits and vegetables (salads) to prevent browning.

### 4-Salts and Sugars

These compounds are grouped together because of the similarity in their modes of action in preserving foods by lower the water activity and have an adverse effect on microorganisms. Sodium chloride is used in brines and curing solutions or applied directly to the food. The early food uses of salt were for the purpose of preserving meats. This use is based on the fact that at high concentrations, salt exerts a drying effect on both food and microorganisms.

Salt has been reported to have the following effects on microorganisms:

1. It causes high osmotic pressure and hence plasmolysis of cells.
2. It dehydrates foods by drawing out and typing up moisture as it dehydrates microbial cells.
3. It ionizes to yield the chlorine ion.
4. It reduces the solubility of oxygen in the moisture
5. It sensitizes the cell against CO<sub>2</sub>.
6. It interferes with the action of proteolytic enzymes.

**Sugar** such as glucose or sucrose has ability to make water unavailable to organisms by osmotic effect.

### **5-H<sub>2</sub>O<sub>2</sub>**

A solution of H<sub>2</sub>O<sub>2</sub> is recommended as an antimicrobial agent in raw milk to be used in cheese processing (to control growth of psychrotrophic Gram negatives that produce heat-stable enzymes), liquid egg to facilitate destruction of *Salmonella* by low-heat pasteurization, packaging material used in aseptic packaging of foods, and food processing equipment. In raw milk and liquid egg, catalase is used before pasteurization to hydrolyze H<sub>2</sub>O<sub>2</sub> to water and oxygen. H<sub>2</sub>O<sub>2</sub> is a strong oxidizing agent, and the germicidal action is associated with this property .

### **6-Epoxides (Ethylene Oxide, Propylene Oxide)**

Ethylene oxide and propylene oxide are used as fumigants to destroy microorganisms (and insects) in grains, cocoa powder, gums, nuts, dried fruits, spices, and packaging materials. They are germicidal and effective against cells, spores, and viruses. Epoxides are alkylating agents and react with various groups (e.g., -SH, -NH, and -OH) in cellular macromolecules, particularly structural proteins and enzymes.

### **7-Antibiotics**

Antibiotics in foods were approved by the Food and Drug Administration (FDA) and approved by the Expert Committee of the World Health Organization (WHO) . These antibiotics include :

**Tetracyclines** to extend the refrigerated shelf life of seafood and poultry . However, because of the possible increase in antibiotic-resistant bacteria, the use of these antibiotics in food was later banned. **Natamycin**, is an antifungal agent. Its use as a dip or spray to prevent growth of molds and formation of mycotoxins on the surface of some cheeses, sausages, and in raw peanuts. **Tylosin**, inhibits protein synthesis, is a bactericidal antibiotic that is more effective against Gram-positive than Gram-negative bacteria and also inhibits outgrowth of germinated

endospores. Because of its high heat resistance, it has been used in controlling the growth of sporeformers in low-acid canned products. **Subtilin** is effective in canned foods, preventing the outgrowth of germinating endospores.

### **7- Diacetyl**

Diacetyl is produced by several species of lactic acid bacteria in large amounts, particularly through the metabolism of citrate. It is antibacterial against many Gram-positive and Gram-negative bacteria. The antibacterial action is probably produced by deactivating some important enzymes.

### **8-Bacteriocins of Lactic Acid Bacteria**

The term **bacteriocin** is currently used to refer to a group of bioactive peptides produced by many bacterial strains from Gram-positive and Gram-negative groups. The bacteriocins produced by many strains of lactic acid bacteria and some propionic acid bacteria are of special interest in food microbiology because of their bactericidal effect normally to different Gram-positive spoilage and pathogenic bacteria and under stressed conditions to different Gram-negative bacteria important in food. Bacteriocins of lactic acid bacteria are bactericidal to sensitive cells, and death occurs very rapidly at a low concentration. Normally, Gram-positive bacterial strains are more sensitive to bacteriocin than Gram negative bacterial strains. Bacteriocins of food-grade lactic acid bacteria, with special reference to nisin and pediocin. Nisin is produced by *Lactococcus lactis*, however, is used in certain canned foods, most often employed in dairy products—processed cheeses, condensed milk, pasteurized milk. Because of the effectiveness of nisin in preventing the outgrowth of germinating endospores of *Cl.botulinum* and this agent has been used as a possible replacement for nitrite in processed meats.

### **9-Spices**

Many spices, condiments, and plant extracts are known to contain antimicrobial compounds. Some of these include cinnamic aldehyde in cinnamon; eugenol in cloves and cinnamon. Their bacteriostatic and fungistatic properties depend on the active agent. Because of the small amounts used as spices in foods, they probably do not produce any antimicrobial effects. However, the antimicrobial components can be used in higher concentrations as oleoresins or essential oils. The antimicrobial properties of garlic, onion, and ginger, as well as cabbage, brussels sprouts, carrots, and others have generated interest for their possible use as natural preservatives.

## **10-Wood Smoke**

Many processed meat products and fishes are processed with smoke generated by burning hardwood. The main reason for smoking meat, fish, and cheese is to impart desirable flavor, texture, and color to the products. The other benefit is the long shelf life of smoked products, especially those exposed to smoke during heating. The smoke contains several different types of chemicals that deposit on the food surface, many of which have antibacterial properties. The most important antibacterial agents are formaldehyde, phenols, and cresols. Depending on the temperature and time of heating, degree of surface drying and the concentrations, smoking can be both bacteriostatic and bactericidal to bacterial cells. Although smoke has a slight antifungal action, it does not have any adverse effects on the survival or germination of bacterial spores. Smoke also contains some chemicals that are carcinogenic, such as benzopyrene and dibenzanthracene.

## **\*Biopreservation**

Biopreservation is the preservation of food by naturally present or artificially introduced microorganisms, or by antimicrobial substances produced by these. The biopreservation of food, especially utilizing lactic acid bacteria (LAB) that are inhibitory to food spoilage microorganisms. Beneficial bacteria or the fermentation products produced by these bacteria are used in biopreservation to control spoilage and pathogens in food. There are a various modes of action through which microorganisms can interfere with the growth of others such as organic acid, hydrogen peroxide, Diacetyl, bacteriocins production ,etc.