

Physical Pharmacy lab

Lab no. 1:

Concentration Expression

Done By:
Lecturer Zeina Dawood

Assistant Lecturer Hiba Sabah



Introduction

The term “physical pharmacy” stems from the application of physical chemistry principles to the area of pharmacy in the design of drug molecules and drug products.



When the physical chemical and biological properties of drug molecules (i.e. preformulation) are understood, it is possible to design dosage forms for selected routes of administration in humans or animals (i.e., formulation). Collectively, the scientific principles applied in the preformulation and formulation processes is termed “physical pharmacy,”

Lab for this course:

Concentration expression,

Two component systems containing liquid phases,

Three component systems,


The tie – line for three component system,

Partition coefficient, and Buffer solutions.


Pharmaceutical mixtures



True solution
(<1 nm) (one phase system)



Colloidal dispersion
(1-500 nm) (as liposome,
microsphere)



Coarse dispersion
(> 500 nm) (as emulsion and
suspension)



➤ **Dispersion :- is pharmaceutical mixture that consist of at least 2phases with one or more dispersed (internal phase)contained in a single continuous (external)phase.**

➤ **Phase :-is defined as the homogeneous physically distinct portion of the system separated from other parts of the system by bounding surface .**

“ True solution will be focused and discussed in this lab. It can be classified according to the states in which the solute and solvent occur, and because 3 states of matter exist, 9 types of homogeneous mixtures of solute and solvent are possible.



Types of Solutions

Solute	Solvent	Example
Gas	Gas	Air
Liquid	Gas	Water in oxygen
Solid	Gas	Iodine vapor in air
Gas	Liquid	Carbonated water
Liquid	Liquid	Alcohol in water
Solid	Liquid	Aqueous sodium chloride solution
Gas	Solid	Hydrogen in palladium
Liquid	Solid	Mineral oil in paraffin
Solid	Solid	Gold—silver mixture, mixture of alloys

Concentration expressions

Expression	Symbol	Definition
Molarity	M, c	Moles (gram molecular weights) of solute in 1 liter of solution
Normality	N	Gram equivalent weights of solute in 1 liter of solution
Molality	m	Moles of solute in 1000 g of solvent
Mole fraction	X, N	Ratio of the moles of one constituent (e.g., the solute) of a solution to the total moles of all constituents (solute and solvent)
Mole percent		Moles of one constituent in 100 moles of the solution; mole percent is obtained by multiplying mole fraction by 100
Percent by weight	% w/w	Grams of solute in 100 g of solution
Percent by volume	% v/v	Milliliters of solute in 100 mL of solution
Percent weight-in-volume	% w/v	Grams of solute in 100 mL of solution
Milligram percent	—	Milligrams of solute in 100 mL of solution



examples

Calculate the Molarity of solution containing 4 gm of NaOH in 500 ml solution (M.wt NaOH =40gm)

$$M = \text{wt.}/\text{M.wt} \quad * \quad 1000/\text{Vol.}(\text{mL})$$

$$M=4/40 * (1000/500) \quad \longrightarrow \quad M= 0.2 \text{ M}$$

Molality

- Calculate Molality of solution containing 2gm NaOH in 50 gm solvent ?
- Molality = $\text{wt} / \text{M.wt} * 1000 / \text{gms of solvent}$
- $m = 2/40 * 1000 / 50$
- $m = 1$

Q/ How many grams of KI (M.wt=166) are needed to prepare 25ml of 0.75 N solution?

$$N = \text{wt} / \text{eq.wt} * 1000 / \text{vol(ml)}$$

$$0.75 = \text{wt} / 166 * 1000 / 25$$

$$\text{wt} = 3.12 \text{ gm}$$

Q/ Solution prepared by dissolving 1 mole ethyl alcohol in 3 moles of water. Calculate mole fraction of each one?

Mole fraction = moles of one constituent (solute) / the total moles of all constituents (solute & solvent)

$$\text{Mole fraction of ethyl alcohol (X}_1\text{)} = 1/1+3 = 1/4 = 0.25$$

$$\text{Mole fraction of water (X}_2\text{)} = 3/1+3 = 3/4 = 0.75$$

★ the sum of mole fraction is equal to one)

■ Mole percent

■ 5%w/w NaOH mean 5 gm NaOH in 100 gm solvent

■ 10%w/v NaOH = 10 gm NaOH in 100ml solvent

■ 15 %v/v NaOH = 15 ml NaOH in 100ml solvent

Q/prepare 25 gm of 4%w/w NaOH?

$$\left[\begin{array}{cc} 4\text{gm} & 100\text{gm} \\ X & 25\text{gm} \end{array} \right] \quad x = 1\text{gm of NaOH with } 24 \text{ gm of water}$$

Q/prepare 25 ml of 4%w/v NaOH?

$$\left[\begin{array}{cc} 4\text{gm} & 100\text{mL} \\ X & 25\text{mL} \end{array} \right]$$

x = 1gm of NaOH put in volumetric flask then complete the volume to 25 mL



Disadvantages of molarity & normality:

1-changing in value with temperature because of expansion or contraction of liquid so can not be use when studying properties of solution at various temp.

2-because Solvent volume in M&N are not really known it is difficult to study properties such as vapor pressure & osmotic pressure which are related to the conc. of solvent.



(because Molality (m) has not the above disadvantages it is used more likely in theoretical studies more than N & M)



Experimental work:

Materials and equipment:

- NaCl, Na₂CO₃, NaOH, Alcohol, Water.
- Volumetric flasks (50cc), pipettes.

Prepare the following solution :

- A. 50 ml of 0.5 M NaCl.
- B . 50 ml of 2N NaCl.
- C. 50 ml of 0.1N Na₂CO₃.
- D. 50 ml of 0.1 M Na₂CO₃.
- E . 50 gm of 2% w/w NaCl solution.
- F. 50 ml of 10% w/v NaOH or NaCl.
- G. 50 ml of 10% v/v alcohol.

M.Wt of NaCl =58.5

M.Wt of Na₂CO₃ = 106

Density = mass/volume

Density of water =1

Example:

- **A. 50 ml of 0.5 M NaCl.**
- $M = \text{wt.}/M.\text{wt} \quad * \quad 1000/ \text{Vol.}(\text{mL})$
- $0.5M = \text{wt.}/58.5 * (1000/50\text{mL})$
- $20\text{wt} = 0.5 * 58.5$
- $\text{wt} = 1.46 \approx 1.5 \text{ gm of NaCl}$
- And so on for another examples

A watercolor illustration featuring the words "Thank you" in a golden-brown, cursive script. The text is surrounded by several pink and purple flowers with dark centers, green leaves, and clusters of small orange and yellow berries. The entire composition is set against a white background.

Thank you