



SAMPLING METHODS

L-2

Prof Dr Najlaa Fawzi

POPULATION:

The largest collection of anything, if this collection has limits, this is finite population, and if not, this is infinite population.

It can be

A-Population of entities: is the largest collection of entities in which we have an interest at a particular time (e.g. population of humans); each population member has many variables.


B-Population of values: it is the largest collection of values of a random variable from which we have an interest of a variable for a particular time e.g. blood urea.

THE POPULATION
All of the individuals of interest

The sample
is selected from
the population

THE SAMPLE
The individuals selected to
participate in the research study

The results
from the sample
are generalized
to the population



SAMPLING

- ▶ A **sample** is “a smaller (but hopefully representative) collection of units from a population used to determine truths about that population” Why sample?
 - ▶ Resources (time, money) and workload
 - ▶ Gives results with known accuracy that can be calculated mathematically

Sampling definition: Sampling is a technique of selecting individual members or a subset of the population to make statistical inferences from them and estimate characteristics of the whole population.

POPULATION

- The measurable quality is called a parameter.
- The population is a complete set.
- Reports are a true representation of opinion.
- It contains all members of a specified group.

SAMPLE

- The measurable quality is called a statistic.
- The sample is a subset of the population.
- Reports have a margin of error and confidence interval.
- It is a subset that represents the entire population.

- ▶ **What is your population of interest?**
 - ▶ To whom do you want to generalize your results?
 - ▶ All doctors
 - ▶ School children
 - ▶ Women aged 15-45 years
 - ▶ Other
- ▶ **Can you sample the entire population?**

SAMPLING

- ▶ **3 factors that influence sample representativeness**
 - ▶ Sampling procedure
 - ▶ Sample size
 - ▶ Participation (response)
- ▶ **When might you sample the entire population?**
 - ▶ When your population is very small
 - ▶ When you have extensive resources
 - ▶ When you don't expect a very high response

The Sampling Design Process

Define the Population



Determine the Sampling Frame



Select Sampling Technique(s)



Determine the Sample Size



Implement the Sampling Process

Process

- ▶ The sampling process comprises several stages:
 - ▶ Defining the population of concern
 - ▶ Specifying a sampling frame, a set of items or events possible to measure
 - ▶ Specifying a sampling method for selecting items or events from the frame
 - ▶ Determining the sample size
 - ▶ Implementing the sampling plan
 - ▶ Sampling and data collecting
 - ▶ Reviewing the sampling process

Population definition

- ▶ A population can be defined as including all people or items with the characteristic one wishes to understand.
- ▶ Because there is very rarely enough time or money to gather information from everyone or everything in a population, the goal becomes finding a representative sample (or subset) of that population.

Population definition

- ▶ Note also that the population from which the sample is drawn may not be the same as the population about which we actually want information. Often there is large but not complete overlap between these two groups due to frame issues etc.
- ▶ Sometimes they may be entirely separate - for instance, we might study rats in order to get a better understanding of human health, or we might study records from people born in 2013 in order to make predictions about people born in 2014.

SAMPLING FRAME

The sampling frame is the list from which the potential respondents are drawn

- Registrar's office
- Class lists
- Must assess sampling frame errors

- *A sampling frame* has the property that we can identify every single element and include any in our sample .
- The sampling frame must be representative of the population

Types of sampling methods: they are probability methods and non-probability methods, the problem in the second type that they cannot be generalized.

Types of Samples

- ▶ **Probability (Random) Samples**
- ▶ Simple random sample
 - ▶ Systematic random sample
 - ▶ Stratified random sample
 - ▶ Multistage sample
 - ▶ Multiphase sample
 - ▶ Cluster sample
- ▶ **Non-Probability Samples**
 - ▶ Convenience sample
 - ▶ Purposive sample
 - ▶ Quota
 - ▶ Snow ball sample

PROBABILITY SAMPLING

- ▶ A probability sampling scheme is one in which every unit in the population has a chance (greater than zero) of being selected in the sample, and this probability can be accurately determined.
- ▶ . When every element in the population does have the same probability of selection, this is known as an 'equal probability of selection' (EPS) design. Such designs are also referred to as 'self-weighting' because all sampled units are given the same weight.


In a population of 1000 members, every member will have a $1/1000$ chance of being selected to be a part of a sample. Probability sampling eliminates bias in the population and gives all members a fair chance to be included in the sample.

PROBABILITY SAMPLING

- ▶ **Simple Random Sampling,**
- ▶ **Systematic Sampling,**
- ▶ **Stratified Random Sampling,**
- ▶ **Cluster Sampling**
- ▶ **Multistage Sampling.**
- ▶ **Multiphase sampling**

SIMPLE RANDOM SAMPLING

Stages in random sampling:



Is also known as '**unrestricted random sampling**'
- Used in clinical trials

SIMPLE RANDOM SAMPLING

- Applicable when population is small, homogeneous & readily available
- All subsets of the frame are given an equal probability. Each element of the frame thus has an equal probability of selection.
- It provides for greatest number of possible samples. This is done by assigning a number to each unit in the sampling frame.
- One of the best probability sampling techniques that helps in saving time and resources
- A table of random number or lottery system is used to determine which units are to be selected.

- ▶ Estimates are easy to calculate.
- ▶ Simple random sampling is always an EPS design, but not all EPS designs are simple random sampling.
- ▶ Disadvantages
 - ▶ It needs complete list of study population , which is often difficult to obtain.
 - ▶ If sampling frame large, this method useless.
 - ▶ Minority subgroups of interest in population may not be present in sample in sufficient numbers for study.

REPLACEMENT OF SELECTED UNITS

- ▶ Sampling schemes may be without replacement ('**WOR**' - no element can be selected more than once in the same sample) or with replacement ('**WR**' - an element may appear multiple times in the one sample).
- ▶ For example, if we catch fish, measure them, and immediately return them to the water before continuing with the sample, this is a WR design, because we might end up catching and measuring the same fish more than once. However, if we do not return the fish to the water (e.g. if we eat the fish), this becomes a WOR design.

Methods of Simple random sampling:

- Lottery method
- Random no. tables
- * Computer software.

The diagram illustrates the lottery method for simple random sampling. It features a black hat with a white brim, labeled 'Hat' at the bottom. Inside the hat, three numbered tickets are visible: a yellow ticket with the number 24, a pink ticket with 38, and a blue ticket with 14. To the right of the hat, a grey ticket with the number 43 is shown. The entire scene is set against a black background with a white title box at the top that reads 'Simple Random Sampling'. Below the title, a pink box labeled 'Method 1' is present. To the right of the hat, there are three pink text boxes: the first contains an example, the second lists the first two steps of the process, and the third lists the final step. A green arrow points from the 'Lottery method' text in the main list to the 'Method 1' box.

Simple Random Sampling

Method 1

Example: Out of a CLASS of 50 students 15 are to be selected to take part in a FEST.

1. Assign ROLL number from 1 to 50 to each student.
2. Write each number on a piece of paper (or use raffle tickets), place in a hat and mix up.
3. Draw the 15 numbers from the hat.

Hat

SYSTEMATIC SAMPLING

- ▶ Systematic sampling depend on arranging the target population according to some ordering scheme and then selecting elements at regular intervals through that ordered list.
- ▶ This method is preferred when the population is large , scattered and not homogenous .


Systematic sampling involves a random start and then proceeds with the selection of every k th element from then onwards.

In this case, $k = (\text{population size} / \text{sample size})$. It is important that the starting point is not automatically the first in the list, but is instead randomly chosen from within the first to the k th element in the list.

Systematic Random Sampling

- Based on sampling fraction: Every K th unit is chosen in the population list, where K is chosen by sampling interval
- Sampling Interval (K) $Q = \text{Total no. of units in population} / \text{Total no. of units in sample}$
- Applicable for large, non-homogenous populations where complete list of individuals is available
- For example, if there is a population of 1000 from which sample of 20 is to be chosen, then $K = 1000/20 = 50$; thus every 50th unit will be included in the sample (i.e. 1st, 51st, 101st, so on...)
First unit among first 50 is chosen by simple random sampling.

SYSTEMATIC SAMPLING.....



As described above, systematic sampling is an EPS method, because all elements have the same probability of selection (in the example given, one in ten). It is *not* 'simple random sampling' because different subsets of the same size have different selection probabilities - e.g. the set $\{4, 14, 24, \dots, 994\}$ has a one-in-ten probability of selection, but the set $\{4, 13, 24, 34, \dots\}$ has zero probability of selection.

▶ ADVANTAGES:

- ▶ Sample easy to select
- ▶ Suitable sampling frame can be identified easily
- ▶ Sample evenly spread over entire reference population
- ▶ Time and labour for sample collection is relatively small.

▶ DISADVANTAGES:

- ▶ Sample may be biased if hidden periodicity in population coincides with that of selection.
- ▶ Difficult to assess precision of estimate from one survey.

STRATIFIED SAMPLING

This method used when the population is not homogenous and is composed of diverse segments.

Where population embraces a number of distinct categories, the frame can be organized into separate "strata." Each stratum is then sampled as an independent sub-population, out of which individual elements can be randomly selected.

Every unit in a stratum has same chance of being selected.

Every unit in a stratum has same chance of being selected. This method gives more representative sampling than simple random sampling in a given large population.

Stratified Random Sampling

- Non-homogenous population is converted to homogenous groups/classes (strata); sample is drawn from each strata at random, in proportion to its size
- Applicable for large non-homogenous population
 - Gives more representative sample than simple random sampling
- None of the categories is under or over-represented
 - For example, In a population of 1000, sample of 100 is to be drawn for Hemoglobin estimation; first convert non-homogenous population is converted to homogenous strata (i.e. 700 males and 300 females), then draw 70 males and 30 females randomly respectively

**Define
population**



**Develop
sampling
frame
according to
characteristics
required**



**Determine
the
proportion of
each
population
variable of
interest**



**Systematic
sampling
methods can
then be
followed to
select sample
unit**

Using same sampling fraction for all strata ensures proportionate representation in the sample.

Adequate representation of minority subgroups of interest can be ensured by stratification & varying sampling fraction between strata as required.

- ▶ Finally, since each stratum is treated as an independent population, different sampling approaches can be applied to different strata.

■ Types of Stratified Samples

■ Proportional Stratified Sample:

- The number of sampling units drawn from each stratum is in proportion to the relative population size of that stratum

■ Disproportional Stratified Sample:

- The number of sampling units drawn from each stratum is allocated according to analytical considerations e.g. as variability increases sample size of stratum should increase

- **Optimal allocation stratified sample**
 - **The number of sampling units drawn from each stratum is determined on the basis of both size and variation.**

 - **Calculated statistically**

Advantage

- It is more representative
- It gives estimates with increased precision
- As the population is more concentrated, the time and money will be saved.

Disadvantage

□ Requires accurate information on proportions of each stratum

It is very difficult task to divide the population into homogenous strata .

□ Stratified lists costly to prepare. This may require considerable time , money and statistical expertise.

Disadvantages to using stratified sampling.

First, sampling frame of entire population has to be prepared separately for each stratum
Second, when examining multiple criteria, stratifying variables may be related to some, but not to others, further complicating the design, and potentially reducing the utility of the strata.

Finally, in some cases (such as designs with a large number of strata, or those with a specified minimum sample size per group), stratified sampling can potentially require a larger sample than would other methods

Select a stratified random sample of 20 patients from 200 patients.

	Disease A	Disease B	Disease C	Disease D	TOTAL
No. of patients	100	60	20	20	200
%	50	30	10	10	100

Out of 20 patients the no. to be selected wise area.

Strata Disease A ($100/200$) X 20 = 50% OF 20 = 10

Disease B ($60/200$) X 20 = 30% OF 20 = 6

Disease C ($20/200$) X 20 = 10% OF 20 = 2

Disease D ($20/200$) X 20 = 10% OF 20 = 2

Cluster Random Sampling

-Applicable when units of population are natural groups or clusters.

* Clusters are heterogeneous within themselves but homogenous with respect to each other

* Clusters are identified and included in a sample based on demographic parameters like age, sex, location, etc.


- Often used to evaluate vaccination coverage in EPI

- ▶ Cluster sampling is an example of 'two-stage sampling' .
- ▶ First stage a sample of areas is chosen;
- ▶ Second stage a sample of respondents *within* those areas is selected.
- ▶ Population divided into clusters of homogeneous units, usually based on geographical contiguity.
- ▶ Sampling units are groups rather than individuals.
- ▶ A sample of such clusters is then selected.
- ▶ All units from the selected clusters are studied.

Two types of cluster sampling methods

One-stage sampling. All of the elements within selected clusters are included in the sample.

Two-stage sampling. A subset of elements within selected clusters are randomly selected for inclusion in the sample.



▶ **Advantages :**

- ▶ Cuts down on the cost of preparing a sampling frame.
 - ▶ This can reduce travel and other administrative costs.
 - ▶ Requires list of all clusters, but only of individuals within chosen clusters
 - ▶ Can estimate characteristics of both cluster and population
-
- ▶ Sampling interval is also calculated in CRS
 - ▶ Accuracy : Low error rate of only $\pm 5\%$

Disadvantages:

Sampling error is higher for a simple random sample of same size. Clusters cannot be compared with each other

Often used to evaluate vaccination coverage in EPI

Use in India: Evaluation of immunization coverage

-- WHO technique used: 30 × 7 technique (total = 210 children)

-- WHO technique used in CRS: 30 × 7 technique (total = 210 children)

* 30 clusters, each containing

* 7 children who are 12 - 23 months age and are completely immunized for primary immunization (till Measles vaccine)

Difference Between Strata and Clusters

- ▶ Although strata and clusters are both non-overlapping subsets of the population, they differ in several ways.
- ▶
- ▶ All strata are represented in the sample; but only a subset of clusters are in the sample.
- ▶ With stratified sampling, the best survey results occur when elements within strata are internally homogeneous.
- ▶ However, with cluster sampling, the best results occur when elements within clusters are internally heterogeneous

Moreover, by avoiding the use of all sample units in all selected clusters, multistage sampling avoids the large, and perhaps unnecessary, costs associated with traditional cluster sampling.

MULTISTAGE SAMPLING

- ▶ Complex form of cluster sampling in which two or more levels of units are embedded one in the other.
- ▶ First stage, random number of districts chosen in all states.
- ▶ Followed by random number of towns , villages.
- ▶ Then third stage units will be houses.
- ▶ All ultimate units (houses, for instance) selected at last step are surveyed.

- ▶ This technique, is essentially the process of taking random samples of preceding random samples.
- ▶ Not as effective as true random sampling, but probably solves more of the problems inherent to random sampling.
- ▶
- ▶ An effective strategy because it banks on multiple randomizations. As such, extremely useful.
- ▶ Multistage sampling used frequently when a complete list of all members of the population not exists and is inappropriate.

Multistage Random Sampling

-Is done in successive stages; each successive sampling unit is nested in the previous sampling unit. For example, in large country surveys, states are chosen, then districts, then villages, then every 10th person in village as final sampling unit.

Advantage:

- Introduces flexibility in sampling. This method is very helpful in many large scale surveys where population list preparation is difficult.
- It is less expensive and less time consuming.
- It permits available resources to be concentrated on limited numbers of units of the frame.

Disadvantages

- ❑ Sampling error is usually increased.
- ❑ Sampling units will be of unequal size at various stages resulting in analytical difficulties.

MULTI PHASE SAMPLING

- **Is done in successive phases**
- ▶ Part of the information collected from whole sample & part from subsample.
- ▶ Study of nutrition ,all the families in the original sample are covered for KAP study in 1st phase. A sub -sample of the families is than surveyed for dietary intake in 2nd phase
- ▶ Then a sub-sample of family members covered in 2nd phase is subjected to anthropometric examination in 3rd phase.
- ▶ Survey by such procedure is less costly, less difficult & more purposeful

A further sub -sample from 3rd phase is subjected to Biochemical tests in 4th phase
Thus the number of subject or units gets reduced in every Succeeding phase, thereby reducing the magnitude of the Complicated and costly procedure reserved for the last Phase.

Survey by such procedure is less costly, less difficult & more purposeful

MATCHED RANDOM SAMPLING

A method of assigning participants to groups in which pairs of participants are first matched on some characteristic and then individually assigned randomly to groups.

- ▶ The Procedure for Matched random sampling can be briefed with the following contexts.
- ▶ Two samples in which the members are clearly paired, or are matched explicitly by the researcher. For example, IQ measurements or pairs of identical twins.

- Those samples in which the same attribute, or variable, is measured twice on each subject, under different circumstances. Commonly called **repeated measures**.
- Examples include the times of a group of athletes for 1500m before and after a week of special training; the milk yields of cows before and after being fed a particular diet.

Uses of probability sampling

There are multiple uses of probability sampling.

They are:

- **Reduce Sample Bias:** Using the probability sampling method, the bias in the sample derived from a population is unimportant to non-existent. The selection of the sample mainly describes the understanding and the inference of the researcher. Probability sampling leads to higher quality data collection as the sample appropriately represents the population.

• **Diverse Population:** When the population is infinite and different, it is essential to have adequate representation so that the data is not tilted towards one demographic. For example, if Square would like to understand the people that could make their point-of-sale devices, a survey conducted from a sample of people across the US from different industries and socio-economic backgrounds helps.

• **Create an Accurate Sample:** Probability sampling helps the researchers plan and create an accurate sample. This helps to obtain well-defined data.

Identify each of the following examples as qualitative) or numerical (quantitative) variables.

- 1. The place of residence for each student in a statistics class.**
- 2. The amount of gasoline pumped by the next 10 customers at the gas station .**
- 3. The amount of water in the tanks of each of 25 homes in certain city.**
- 4. The color of the T-shirts worn by each of 20 children .**
- 5. The length of time to complete a statistic homework assignment.**

Example: Identify each of the following as examples of (1) nominal, (2) ordinal, (3) discrete, or (4) continuous variables:

- 1. The length of time until a pain reliever begins to work.**
- 2. The number of chocolate chips in a cookie.**
- 3. The number of colors used in a statistics textbook.**
- 4. The brand of refrigerator in a home.**
- 5. The overall satisfaction rating of a new car.**
- 6. The number of files on a computer's hard disk.**
- 7. The pH level of the water in a swimming pool.**
- 8. The number of doctors in a health center .**

END OF PART ONE