

**By the Name of ALLAH the Most Gracious the Most Merciful**



# The peritoneum, mesentery, greater omentum and retroperitoneal space Part II

د . أحمد اسامة حسن

**Specialist in General Surgery & Laparoscopic Surgery**

**28<sup>th</sup> ,September , 2023**

To be read in

**Bailey & Love's Short Practice of Surgery 27<sup>th</sup> Edition.**

**Ch 65.**

# Part II

- **Abdominal Fluid Collections :**

- Intraperitoneal collections ( Ascites ).
- Retroperitoneal collections.

- **MESENTERY**

MESENTERIC HAEMATOMA

MESENTERIC ADENITIS

IN CROHN'S DISEASE

TUBERCULOSIS OF THE MESENTERIC LYMPH NODES

Mesenteric panniculitis

Adhesions ( types , Mx, Cx ).

VASCULAR ABNORMALITIES OF THE MESENTERY

Mesenteric cysts (Chylolymphatic cysts, Enterogenous ).

# Abdominal Fluid Collections

- Subdivided into:
- Intraperitoneal collections ( Ascites )
- Retroperitoneal collections. are further subdivided into :
  - Those limited to the retroperitoneal space (e.g. in pancreatitis)
  - Collections arising in relation to retroperitoneal organs such the kidney.

# Ascites

# Causes of Ascites

## Transudates (protein <25 g/L)

- Low plasma protein Concentrations
  - Malnutrition
  - Nephrotic syndrome
  - Protein-losing enteropathy
  - Cirrhosis
- High central venous pressure
  - Congestive cardiac failure
  - Constrictive pericarditis  
(Pick's disease)
  - Vena cava obstruction
- Portal hypertension
  - Portal vein thrombosis
  - Cirrhosis .
  - Budd–Chiari syndrome ( Chronic )  
(hepatic vein occlusion or thrombosis)

## Exudates (protein >25 g/L)

- Peritoneal malignancy
- Bacterial peritonitis.
- Tuberculous peritonitis
- Budd–Chiari syndrome ( Acute )  
(hepatic vein occlusion or thrombosis)
- Pancreatic ascites
- Chylous ascites
- Meigs' syndrome ( reversible )

## Ascites : Accumulation of peritoneal fluid, if there is excess production or reduced absorption.

- Peritoneal fluid is constantly secreted and absorbed.
- Production of large volumes of a protein-rich fluid occurs in peritonitis and carcinomatosis peritonei.
- Reduced absorption ( capillary pressure is increased as a result of generalised water retention, cardiac failure, constrictive pericarditis or vena cava obstruction. Capillary pressure is also raised selectively in the portal venous system in the Budd–Chiari syndrome, hepatic cirrhosis or extrahepatic portal venous obstruction.
- Plasma colloid osmotic pressure may be lowered in patients with malnutrition, diminished intestinal absorption, abnormal protein losses or defective protein synthesis ( cirrhosis)
- Peritoneal lymphatic drainage may be impaired, resulting in the accumulation of protein-rich fluid ( metastasis ).

# Examination :

- Underlying disease.
- Becomes clinically recognizable when greater than 1.5 liters of fluid is apparent (although greater volumes may be required in obese patients).
- The abdomen is distended evenly with fullness of the flanks, tense, which are dull to percussion.
- Usually, shifting dullness is present.
- But, when there is a very large accumulation of fluid, this sign is absent. In such cases, flicking the abdominal wall produces a characteristic transmitted fluid thrill on the other side of the abdomen. This is not a reliable clinical sign. In women, ascites must be differentiated from an enormous ovarian cyst.

# Investigations

- Investigation for the cause.
- Liver function tests (LFTs), cardiac function, ultrasonography and/or CT scanning .
- Ascitic aspiration or tap under imaging guidance ( Dx & Tx ) . sent for microscopy/cytology, culture, including mycobacteria & and analysis of protein content and amylase.



# Management

- Therapeutic / Palliative.
- Portal hypertension : TIPSS.
- Dietary sodium restriction to 200 mg/day may be helpful, but diuretics are usually required (combination of spironolactone and furosemide).
- Paracentesis can be performed. Serial large volume paracentesis (4–6 L/day and up to 8 litres in one session) can be performed safely with albumin replacement and can be performed in patients with cirrhosis and deranged clotting.
- Indwelling external drain for smaller volume home paracentesis.

# Chylous ascites

- The ascitic fluid appears milky because of an excess of chylomicrons (triglycerides).
- Most of these cases are associated with malignancy (usually lymphoma).
- Other causes include cirrhosis, TB, filariasis, nephrotic syndrome, abdominal trauma (including surgery), constrictive pericarditis, sarcoidosis and congenital lymphatic abnormality.
- The prognosis is poor unless the underlying condition can be cured.
- In addition to other measures used to treat ascites, patients should be placed on a fat-free diet with medium-chain triglyceride supplements.

# The Mesentery

# GENERAL CONSIDERATIONS

- The mesentery comprises adipose, connective tissue, neurological, lymphatic and vascular components. Abnormalities can arise in any of these and lead to either solid (tumour deposits, lymphatic metastases) or cystic lesions.
- The supportive capacity of the mesentery is reflected in the finding of splenunculi, heterotopic pancreas, ossification, teratomas and even ectopic pregnancies in different regions of the mesentery

# MESENTERIC HAEMATOMA

can follow

- Abdominal compression in trauma (e.g. seat-belt syndrome)
- During abdominal surgery, when the mesentery must be manipulated.
- A haematoma may form and quickly enlarge to compress mesenteric veins.

# MESENTERIC ADENITIS

. It mostly occurs in the ileocaecal region because of the volume of lymphatic tissue.

It is often the site of viral or infective lymphadenopathy (Yersinia spp., Campylobacter spp., Mycobacterium tuberculosis) and may follow an upper respiratory tract infection with either a viral or bacterial pathogen.

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is not uncommon.

The patient is frequently pyretic (often the temperature is markedly elevated) and may have enlarged cervical lymph nodes.


- During childhood, acute, non-specific mesenteric adenitis is a common condition. The typical history is one of short attacks of central abdominal pain lasting from 10 to 30 minutes, commonly associated with vomiting. The patient seldom looks ill. In more than half of the cases the temperature is elevated. Abdominal tenderness is poorly localised and, when present, shifting tenderness ( move to other side ) is a valuable sign for differentiating the condition from appendicitis. The neck, axillae and groins should be palpated for enlarged lymph nodes.

(Looking for tonsillitis ).

**DDx : acute appendicitis**

Right,  
mesocolon.

Mesenteric  
adenitis.





## Investigation

- There is often a leukocytosis of 10 000–12 000/ $\mu$ L or more on the first day of the attack, but this falls on the second day.
- Ultrasonography may be helpful in differentiating this from appendicitis.
- Sometimes a CT or exploratory laparoscopy is required.

## Treatment


- This is normally supportive.
- Viral mesenteric adenitis normally resolves spontaneously but can recur.
- In bacterial mesenteric adenitis include cramping pain, vomiting and diarrhoea. They can be severe and require hospitalisation.

# MESENTERIC ADENITIS AND THE MESENTERY IN CROHN'S DISEASE

- Ileocaecal mesenteric adenitis occurs in ileocolic Crohn's disease and the mesentery is thickened, shortened and oedematous with a tendency to bleed if handled. The mesenteric vascular pedicles may not be apparent, thus great care is needed when dividing the mesentery as normal techniques may not be suitable.
- Specifically the mesentery can extend over adjoining intestine as 'fat wrapping' or 'creeping fat' .
- There is mesenteric transition zone. ( surgical excision would be at its intersection with adjoining mesentery ).

Mesenteric  
transition zone

Fat wrapping  
Creeping fat



Normal  
mesentery

# TUBERCULOSIS OF THE MESENTERIC LYMPH NODES

- Is considerably less common than acute non-specific lymphadenitis.
- Usually, but not necessarily, bovine in origin, are ingested and enter the mesenteric lymph nodes by way of Peyer's patches.
- Usually there are several, and occasionally massive involvement occurs. May be Calcified that can be demonstrated on a plain radiograph of the abdomen, DDX ( renal or ureteric stones ) .
- The presentation may be with abdominal pain (a rare differential for appendicitis) or with general constitutional symptoms (pyrexia, weight loss, etc.).
- Ix : CBP, ESR,CRP, CXR, Sputum ( C+S ).
- Rx : Anti TB.

# Mesenteric panniculitis


- This is inflammation of the mesodermal mesentery (i.e. the mesenteric stroma).
- It is always present in Crohn's disease (**Mesenteric adenitis and the mesentery in Crohn's disease**).
- Often it is an incidental finding on cross-sectional imaging of the abdomen by CT ('misty mesentery' ( fat stranding ) ).
- It can arise secondary to inflammation in any digestive organ. It is associated with connective tissue disorders (including Weber– Christian disease).
- Treatment is normally medical and surgery is rarely required. Serial CT scanning is indicated to ensure resolution.

# Adhesions

- Peritoneal (sclerotic) adhesions
- Areolar adhesions.
- Dense adhesions.

# Peritoneal (sclerotic) adhesions

- These are mesothelial adhesions between two mesothelial surfaces .They reflect mesothelial proliferation, resemble peritoneum and are generally soft and non-vascular.
- These can be band-like linking a viscus . They may occur following laparoscopic surgery, .
- Band adhesions can lead to focal abdominal pain, internal herniation (through the window created with surrounding related structures) or intestinal torsion around the band.





# Areolar adhesions

- These flimsy connective tissue adhesions are identical to the connective tissue that fills the retroperitoneal space between the mesentery and posterior abdominal wall .
- Given this appearance they have been called ‘angel hairs’.
- In the most extremes cases, the peritoneal cavity may be obliterated or limited to small pockets at the flanks. This generates considerable technical challenges during reoperative surgery.



## Dense adhesions

- Following severe intraperitoneal contamination (e.g. after perforation)
- They can bridge the abdominal wall and intestine, or intestine and mesentery, and lead to fusion of the bridged structures.
- Surgical division can be challenging. Sometimes it is not possible to separate conjoined organs without disrupting the integrity of one of them.

# Complications of adhesions

- The most is small bowel obstruction (SBO). (Chapter 74).
- As a major cause of secondary infertility (Chapter 87).
- Treatment : Guided division of adhesions ( adhesolysis )
- To date, no agent or mechanism has been identified that reliably reduces adhesion formation.

# Management in SBO

- Conservative.
- Adhesolysis.

# VASCULAR ABNORMALITIES OF THE MESENTERY

- Acute mesenteric ischaemia .
- Chronic mesenteric ischaemia
- Venous ischaemia

# Acute Mesenteric Ischaemia

- Three major vessels: the coeliac trunk and the superior and inferior mesenteric arteries.
- However, anastomoses between peripheral branches of the SMA and IMA, referred to as the marginal artery of Drummond, usually prevent critical ischaemia of the sigmoid and descending colon.
- Embolism : leads to ischaemia and necrosis of most of the intestine.
- At first, the severity of abdominal pain does not match clinical findings on examination. If ischaemia and necrosis occur, the patient develops peritonism as a result of irritation of the parietal peritoneum by the necrotic intestine.
- If ischaemia is limited to the mucosa, the patient may experience cramping suprapubic pain and diarrhoea that normally settles. If ischaemia is transmural the colon may become necrotic and require resection.

# Chronic Mesenteric Ischaemia

- Is due to atherosclerotic narrowing at the origin of any of the three arterial trunks.
- Presentation : postprandial abdominal pain that can be severe, resulting in a fear of eating with progressive weight loss.
- The diagnosis requires CT angiography. Radiological stent placement may be successful, but surgical endarterectomy or bypass repair may be required.



# Venous Ischaemia

- The venous drainage of the mesenteric domain is limited to the junction between the hepatic veins and the inferior vena cava (portal vein) organs , so an alternative drainage route opens (i.e. a portosystemic shunt) or is created .
- Narrowing or blockage of the lumen at this junction occurs in Budd–Chiari syndrome. It has major clinical implications.
- Portal venous thrombosis impedes venous drainage of all abdominal digestive organs unless an alternative drainage route opens (i.e. a portosystemic shunt) or is created

# ROTATIONAL DISORDERS

- Malrotation .
- Volvulus of the intestine and adjoining mesentery .

# Mesenteric cysts

- They are most often observed in mesentery adjoining the small intestine (60%) or the colon (40%) and can be classified as follows:
- Chylolymphatic.
- Enterogenous.
- Traumatic.
- Hydatid.

- Most commonly in adults with a mean age of 45 years
- Twice as common in women as in men .
- Approximately a third of cases occur in children younger than 15 years.
- The mean age of children affected is 5 years.
- Common presentation is of a painless fluctuant abdominal swelling near the umbilicus
- Other presentation recurrent attacks of abdominal pain with or without vomiting (pain is due to recurring temporary impaction of a food bolus in a segment of bowel narrowed by the cyst or possibly from torsion of the mesentery) and acute abdominal catastrophe is due to:
  - Torsion of the related mesentery.
  - Rupture of the cyst .
  - Haemorrhage into the cyst.
  - Infection

# Chylolymphatic cysts

- This is the most common variety arising in congenitally misplaced lymphatic tissue that has no efferent communication with the lymphatic system (most frequently in the mesentery of the ileum).
- The cyst is filled with clear lymph or, less frequently, with chyle, ( watered milk to cream).
- More often unilocular than multilocular, a chylolymphatic cyst is almost invariably solitary.
- It has independent blood supply thus, enucleation is possible without the need for resection of the gut.

# Enterogenous cysts

- These are believed to be derived either from :
- A diverticulum of the mesenteric border of the intestine that has become sequestered from the intestinal canal during embryonic life.
  - Duplication of the intestine .
  - It has a thicker wall than a chylolymphatic cyst and is lined by ciliated mucous membrane. The content is mucinous and either colourless or yellowish brown as a result of past haemorrhage.
  - It have a common blood supply; consequently, removal of the cyst always entails resection of the related portion of intestine.

حَمْدُ اللَّهِ

**PRAISE BE TO ALLAH**