

# **JAUNDICE IN FAMILY MEDICINE**



**Assistant Prof.  
Dr Alaa A.Salih –FICMS(FM) -2026**

Jaundice, is a yellowish discoloration of tissue resulting from the deposition of bilirubin.

Tissue deposition of Bilirubin occurs only in the presence of serum hyperbilirubinemia .



- Increases in serum Bilirubin level are best detected by examining the sclera, which have a particular affinity for bilirubin due to their high elastin content.
- The presence of scleral jaundice indicates a serum bilirubin level of at least 51  $\mu\text{mol/L}$  (3 mg/dL).
- If the examiner suspects scleral jaundice, a second site to examine is underneath the tongue.

- As serum bilirubin levels rise, the skin will eventually become yellow in light-skinned patients and even green
- If the process is long-standing; the green color is produced by oxidation of bilirubin to biliverdin.



# Critical Questions in the Evaluation of the Jaundiced Patient

- Is it jaundice?
- Acute vs. Chronic Liver Disease=duration
- Prehepatic VS hepatic
- Hepatocellular vs. Cholestatic

# Evaluation of the Jaundiced Patient HISTORY

- Pain
- Fever
- Confusion
- Weight loss
- STI
- Alcohol
- Medications
- Itching
- malaise, myalgias
- dark urine
- Abdominal girth
- Edema
- Autoimmune diseases
- HIV status
- Family history liver dis

# Evaluation of the Jaundiced Patient

## EXAMINATION

- Vital signs
- Mental status
- Asterixis
- Abd tenderness
- hepatomegaly
- Splenomegaly
- Ascites
- Edema
- Spider angiomas
- Hyperpigmentation
- Kayser-Fleischer rings
- Xanthomas
- Gynecomastia
- Left supraclavicular adenopathy (Virchow's node)

# Evaluation of the Jaundiced Patient

## LAB EVALUATION

- AST(S.GOT)-ALT(S.GPT)-ALP
- Bilirubin –total/indirect
- Albumin=chronic
- INR&PT= acute
- B.Glucose
- Na-K-PO<sub>4</sub>, acid-base
- Acetaminophen level
- CBC/plt
- Viral serologies
- ANA-ASMA-AMA
- Ceruloplasmin
- Iron profile
- Blood cultures

# Evaluation of the Jaundiced Patient

- Ultrasound:
  - More sensitive than CT for gallbladder stones
  - Equally sensitive for dilated ducts
  - Portable, cheap, no radiation, no IV contrast
- CT:
  - Better imaging of the pancreas and abdomen
- MRCP:
  - Imaging of biliary tree comparable to ERCP
- ERCP:
  - Therapeutic intervention for stones
  - Brushing and biopsy for malignancy

# Case 1

- **A 19-year-old male presents with progressive yellowing of the eyes for 4 days accompanied by fatigue and mild shortness of breath. He reports passing dark urine but denies pale stools, abdominal pain, fever, or weight loss. He had a recent upper respiratory infection one week ago. No history of alcohol use or drug intake.**
- **Examination:**
- **Mild scleral icterus**
- **Pallor present**
- **No hepatomegaly**
- **Mild splenomegaly**
- **Vitals stable**

# Laboratory Investigations:

Test	Result	Reference
Hemoglobin	8.6 g/dL	Low
Reticulocyte count	7%	High
Total bilirubin	4.2 mg/dL	High
Indirect bilirubin	3.5 mg/dL	High
Direct bilirubin	0.7 mg/dL	Mild
AST/ALT	Normal	
ALP	Normal	
LDH	Elevated	
Haptoglobin	Very low	
Peripheral smear	Spherocytes seen	
Direct Coombs test	Positive	

**Most Likely Diagnosis:  
Autoimmune hemolytic anemia causing pre-hepatic jaundice.**

**Teaching Points:**

- **Predominantly unconjugated bilirubin**
- **Normal liver enzymes**
- **Evidence of increased red cell destruction**

# Case 2

**A 52-year-old female presents with yellow discoloration of skin for one-week, intense generalized itching, pale stools, and tea-colored urine. She complains of intermittent right upper quadrant pain, especially after fatty meals. No prior liver disease.**

- **Examination:**
- **Deep jaundice**
- **Scratch marks over limbs**
- **Right upper quadrant tenderness**
- **Afebrile**

# Laboratory Investigations:

Test	Result	Reference
Total bilirubin	9.8 mg/dL	High
Direct bilirubin	7.6 mg/dL	High
Indirect bilirubin	2.2 mg/dL	Mild
ALP	Markedly elevated (620 IU/L)	
GGT	Elevated	
AST/ALT	Mild elevation	
CBC	Normal	
INR	Slightly prolonged	

# Imaging:

- **Ultrasound abdomen: Dilated common bile duct (10 mm) with suspected distal stone; gallbladder contains multiple calculi.**
- **MRCP: Single obstructing stone in the distal common bile duct.**

**Most Likely Diagnosis:  
Obstructive jaundice secondary to common  
bile duct stone.**

**Teaching Points:**

- **Predominantly conjugated bilirubin**
- **Cholestatic enzyme pattern (high ALP, GGT)**
- **Pale stool and pruritus strongly suggest obstruction**

# Clinical Aspects of Jaundice

- Clinically detectable if SB is  $>2.5$  mg%
- With edema and dark skin – Jaundice is masked
- What is special about the sclera ? – Rich Elastin
- Darkening of the urine – Differential Diagnosis
- Skin discoloration – Yellowish, - Carotinemia – Eyes N
- Mucosa – hard palate (in dark skinned)
- Greenish of skin and sclera - due Biliverdin – indicates long standing jaundice
- Generalized Pruritus – Obstructive Jaundice – Why ?

# Clinical History – Imp. clues

- Duration of jaundice – Acute / Chronic
- Abdominal pain v/s painless jaundice
- Fever – Viral / bacteria /sepsis
- Arthralgia, rash, glands; Pruritus - obstructive
- Appetite – Hepatocellular / Malignancy
- Weight loss – Malignancy
- Colour of stools –chalky white –obstructive
- Family history – Hemolytic – Inherited dis.
- H/o transfusion, STI
- Alcohol abuse, Medications – INH, EM,  
Largactil=Chlorpromazine

SUUDEN DISTURBE CONSCIOUSNESS	BP=129/71		WBC=13.6	DIAMICRON 60MG ONCE
HICCOUGH-	HR=87		HB=12.8	BETALOC 100MG
ON LARGACTILE 50MG *2	CANNOT MOVE LT HAND		MCV=62	ASPIRIN 81MG
	NDR		RBS=146/ TSB=5.3	XIGDUO 5/1000*2
	CLEAR CHEST		AST=404/ALT=611	LEGALON
	NO LEG OEDEMA		U=72/HBA1C=6.3	LINCOCIN 600MG=5
	JAUNDICE		CRP=113	BELIRELAX/B12=5/

25/3/2023

HICCOUGH	BP=129/71	U/S=IBS	HB=12.4/MCV=60	LYRICA 75MG
HYPOGLYCEMIA	HR=87		ALT=8.5/AST=9.4	BETALOC 100MG
TO RE-EVALUATE KIVER FUNCTION'	CANNOT MOVE LT HAND		U=50/C=0.5	ASPIRIN 81MG
	NDR		LDL=65	XIGDUO 5/1000*1
	CLEAR CHEST		CRP=17	DUPHALC SYP
	NO LEG OEDEMA		T4=100/TSH=1.65	MAXIM 400MG
HICCOUGH				

DRAASFICMSFM

# Colored Urine – Differ. Diagnosis

- Concentrated urine in dehydration
- Fluid deprivation syndromes
- Sulfasalazine use – for Ulcerative colitis
- Rifampicin, Pyridium and Thiamine use
- Red urine – Porphyria,
- Hemoglobin & Myoglobinuria, Hematuria
- Melanin excretion from Melanoma
- Red sweat in Clofazamine, Rifampicin

# An Approach to Jaundice

- Is it isolated elevation of serum bilirubin ?
- If so, is the ↑ unconjugated or conjugated fraction?
- Is it accompanied by other liver test abnormalities ?
- Is the disorder hepatocellular or cholestatic?
- If cholestatic, is it intra- or extrahepatic?
- These can be answered with a thoughtful History and physical examination
- Interpretation of laboratory tests and Radiological tests and procedures.

# Normal values for LFT

Features	Healthy Normal
Total Bilirubin	<b>0.3-1.3 mg/dl</b>
Conjugated Bilirubin	<b>0.1-0.4 mg</b>
AST (S.GOT)	<b>12-38 U/L</b>
ALT (S.GPT)	<b>7-41 U/L</b>
Alkaline phosphatase	<b>13-100 U /L</b>
GGT and 5' Nucleosidase.	Significantly ↑ in ALD
Urine Bilirubin	Absent
Urine Urobilinogen	In trace quantity
Urine Bile Salts	Absent

# Lab Diagnosis of Jaundice – D.D

Features	Prehepatic (Hemolytic)	Intrahepatic (Hepatocellular)	Posthepatic (Obstructive)
Unconjugated	↑	Normal	Normal
Conjugated	Normal	↑	↑
AST or ALT	Normal	↑ ↑	Normal
Alkaline phos. and GGT	Normal	Normal	↑ ↑
Urine bilirubin	Absent	Present	Increased
Urobilinogen	Increased	Present	Absent
Hb and Retic	Low+ increase	normal	normal

# Utility of Liver Function Tests

<b>LFT</b>	<b>Utility of the test</b>
<b>ALT/SGPT</b>	<b>ALT ↓ than AST in alcoholism</b>
<b>Albumin</b>	<b>Assess severity / chronicity</b>
<b>Alk. phosphatase</b>	<b>Cholestasis, hepatic infiltrations</b>
<b>AST/SGOT</b>	<b>Early Dx. of Liver disease, Follow up</b>
<b>Bilirubin (Total) /Conjug.</b>	<b>Diagnose jaundice</b>
<b>Gamma-globulin</b>	<b>Dx. F/up Chronic hepatitis &amp; cirrhosis</b>
<b>GGT</b>	<b>Dx alcohol abuse</b>

# Non-Hepatic causes of abnormal LFT

<b>ABNORMAL LFT</b>	<b>NON-HEPATIC CAUSES</b>
<b>Albumin</b>	<b>Nephrotic syndrome Malnutrition, CHF</b>
<b>ALP</b>	<b>Bone disease, Pregnancy, Malignancy , Adv age</b>
<b>AST (S.GOT)</b>	<b>MI, Myositis, I.M.injections</b>
<b>Bilirubin</b>	<b>Hemolysis, Sepsis, Ineffective erythropoiesis</b>
<b>PTT</b>	<b>Antibiotics, Anticoagulant, Steatorrhea, Dietary</b>

# **ALGORITHMIC APPROACH FOR JAUNDICE**

**How to clinically evaluate the patient ?**

**What tests will help us in D.D ?**

**What imaging modalities will be useful ?**

**How to monitor the progress ?**

# First Step

**Estimate Serum Bilirubin**

**Is it less than 1 mg % - Normal**

**Is it more than 1 mg % - Elevated**

# Second Step : If SB > 1.0 mg

Is it unconjugated bilirubin ?

Haemolytic Jaundice

Is it Conjugated Bilirubin ? (> 20%)

Hepatocellular jaundice

Obstructive jaundice

# ↑ in Unconjugated Bilirubin

**Hemolytic Jaundice - Uncommon**

**1. Hemolytic Disorders + Anemia**

**Inherited – Sphero, SS, G6PD, PK**

**Acquired – Paroxysmal nocturnal  
hemoglobinuria**

**2. Ineffective Erythropoiesis –B<sub>12</sub>, Fe, F**

**3. Drugs – Rifampicin, Probenecid**

**4. Inherited –Crigler Najjar, Gilberts**

# Third Step : If Conj SB increased

Do - AST and ALT (SGOT and SGPT)

Elevated AST and ALT

Hepatocellular jaundice

AKP, 5N, GGT will be normal

Do - Alkaline Phosphatase and GGT

AKP, GGT ↑↑ in Obstructive Jaundice

AST and ALT will be normal

# Fourth Step : Hepatocellular

Hepatocellular – Features and D.D

Conjugated SB is increased

AST and ALT are increased

AKP, 5NS, GGT are normal

Hepatitis – A, B, C, D, E, CMV, EBV

Toxic Hepatitis – Drugs, Alcohol

Malignancy – Primary Ca

Cirrhosis – ALD, NAFLD

# What imaging we need

- **Ultrasonography – 98% Sp, 90% Sen.**
- **For GB stones USG better than CT**
- **For duct stones –only 40% seen in USG**
- **PTC – Extrahepatic obstr. – drainage**
- **ERCP – Distal biliary obstruction Dx.Rx.**
- **MRCP – Most useful for duct stones**


# Chronic Liver Disease (CLD)

- **Alcoholic Liver (ALD)**
- **Chronic viral hepatitis**
  - **Hepatitis B**
  - **Hepatitis C**
- **Autoimmune liver disease:**
  - **Autoimmune hepatitis**
  - **Primary Biliary Cirrhosis (PBC)**
- **Inherited conditions**
  - **Haemochromatosis**
  - **Wilson's Disease**
  - **Alpha1-Antitrypsin Deficiency (AATD)**
- **Non-alcoholic fatty liver disease (NAFLD)**
- **Budd-Chiari syndrome**
- **Cryptogenic**

# Hepatotoxic drugs

<b>Conventional Drugs</b>	<b>Natural Substances</b>
<b>Acetaminophen, Alpha-methyldopa</b>	<b>Vitamins, Hypervitaminosis A</b>
<b>Amiodarone, Dantrolene, Diclofenac</b>	<b>Niacin, Cocaine, Mushrooms</b>
<b>Disulfiram, Fluconazole, Glipizide</b>	<b>Aflatoxins, Herbal remedies</b>
<b>Glyburide, Isoniazid, Ketaconazole</b>	<b>Senecio, crotalaria,</b>
<b>Labetalol, Lovastatin, Nitrofurantoin</b>	<b>Pennyroyal oil, Chapparral,</b>
<b>Thiouracil, Troglitazone, Trazadone</b>	<b>Germander, Senna, Herbal mix.</b>

# Acute Cholecystitis



**GB wall is thickened and striated.  
Courtesy of Udo Schmiedl, M.D.**

# Causes of Cholestatic Jaundice

<b>Intrahepatic</b>	<b>Extrahepatic</b>
<b>Acute liver injury, Viral hepatitis</b>	<b>Choledocholithiasis</b>
<b>Alcohol hepatitis, Drugs</b>	<b>Stone obstructing CBD, CD</b>
<b>Chronic liver injury, PBC, PSC</b>	<b>Biliary strictures</b>
<b>Autoimmune cholangiopathy</b>	<b>Cholangiocarcinoma</b>
<b>Drugs, Total parenteral nutrition</b>	<b>Pancreatic carcinoma</b>
<b>Systemic infection, Postoperative</b>	<b>Pancreatitis, Periapillary Ca</b>
<b>Benign causes, Amyloid, lymphoma</b>	<b>PSC, Biliary atresia, duct cysts</b>

# Drugs causing Cholestasis


- **Anabolic steroids (testosterone)**
- **Antithyroid agents (methimazole)**
- **Azathioprine (Immunosuppressive drug)**
- **Chlorpromazine HCl (Largactil)**
- **Clofibrate, Erythromycin estolate**
- **Oral contraceptives (containing estrogens)**
- **Oral hypoglycemics (especially chlorpropamide)**

# Complications of CLD

- **Portal hypertension**
  - **Varices**
  - **Ascites**
  - **Hypersplenism**
- **Synthetic dysfunction**
  - **Coagulopathy**
  - **Encephalopathy**
- **Immunodeficiency**
- **Malnutrition**
- **Hepato-cellular carcinoma**




# Magnetic Resonance Cholangio- Pancreatography (MRCP)



Two stones in the common bile duct


Courtesy of Udo Schmiedl, M.D.

# Retrograde Cholangiogram - ERCP



**Irregular dilation of intrahepatic and extrahepatic ducts.  
Courtesy of Charles Rohrmann, M.D.**

# CT Abdomen



**A large mass with a hepatoma.  
Courtesy of Udo Schmiedl, M.D.**

# When to refer to GE Specialist

**Unexplained jaundice**

**Suspected biliary obstruction**

**Acute hepatitis - severe or fulminant**

**Unexplained abnormal LFTs persisting (for 6 months or greater)**

**Unexplained cholestatic liver disease**

**Cirrhosis (in non-alcoholic) for consideration of liver transplant**

**Suspected hereditary hemochromatosis**

**Suspected Wilson's disease**

**Suspected autoimmune hepatitis**

**Chronic hepatitis C for consideration of antiviral therapy**

**THANK YOU**

---