

Dilution Problems

$V \uparrow M \downarrow$

400 mL
1 M

$$M_1 V_1 = M_2 V_2$$

**Dilutions and
concentrations
Lab 6**

2 M

200 mL

$$= M_3 V_3$$

remove
H₂O

4 M

100 mL


$$M = \frac{\text{mol}}{\text{L}}$$

$V \downarrow M \uparrow$


Dilution and Concentration

- Dilution means when a given solution of a mixture of high concentration is diluted by addition of the suitable diluents or admixture with solution of lower concentration.
- While concentration means when a given solution of a mixture of low concentration are concentrated either by addition of active ingredient or by admixture with higher strength solution or by evaporation of the diluents.
- We have different types of dilution either of liquids or solids.

before




lower volume




higher concentration

Dilution
 $c_1 \cdot V_1 = c_2 \cdot V_2$

after adding solvent




higher volume



lower concentration

Dilution Law

- $C_1 V_1 = C_2 V_2$
- The concentration is expressed either by normality, molarity or percent (%).
- Normality is an expression of the concentration of the solution in terms of equivalent per liter of solution (number of gram eq.wt per 1000ml).
- Molarity is the concentration of the solution in terms of moles per liter.

A hand in a black glove is pouring a clear liquid from a small glass beaker into a larger Erlenmeyer flask. The scene is set against a blue background with other laboratory glassware like a round-bottom flask, a graduated cylinder, and a beaker. A white circular graphic element is overlaid on the right side of the image, containing text.

Addition of extra solvent to the existing solution, in order to reduce its concentration is called 'dilution'.

NOTES :

We have two rules wherever they may be applied will simplify the calculation :

- When ratio strength are given, convert them to % before setting.

Ex: $1:10=10\%$

- Wherever proportional parts enter into calculation, reduce them to the lowest terms.
- Ex: $75:25$ simplify to $3:1$

Dilution of alcohol

- When alcohol is diluted with water a noticeable contraction in volume occurs so it is difficult to calculate the amount of water to be added because alcohol interaction with water by bonding (H-bond) and lead to contraction but this contraction of volume not affect the weight of alcohol and water added.

- Examples

Rx	Boric acid	10gm
	Alcohol 70%	30ml
	Alcohol available	90%

- How many mls of 20% alcohol can be used to prepare 25ml of 10% alcohol?

$$C_1V_1=C_2V_2$$

$$20\% \times V_1 = 10\% \times 25\text{ml}$$

$V_1 = 12.5\text{ml}$ of 20% alcohol and complete the volume to 25ml.

Problems

- If 500ml of 15% v/v solution are diluted to 1500ml. What will be the percentage strength?

$$C_1V_1 = C_2V_2$$

$$15\% \times 500\text{ml} = C_2 \times 1500\text{ml}$$

$$C_2 = 5\%$$

- How many mls of a 1:5000 (w/v) solution of potassium permanganate can be made from 50 ml of a 5% solution?

$$1:5000 = 0.02\%$$

$$C_1V_1 = C_2V_2$$

$$50\text{ml} \times 5\% = 0.02\% \times V_2$$

$$V_2 = 1250\text{ml}$$

Problems

- How much water should be mixed with 5000ml of 85% alcohol to make 50% (v/v) solution?

$$C_1V_1 = C_2V_2$$

$$5000\text{ml} \times 85\% = 50\% \times V_2$$

$$V_2 = 8500\text{ml}$$

$$8500 - 5000 = 3500\text{ml of H}_2\text{O}.$$

- Note
- Standard solution :is a solution of known concentration (normality, molarity and molality) or it's concentration is exactly measured.
- Standardization :is determination of the molarity or normality of the solution.

Reducing and Enlarging Formula

- Pharmacist may have to **reduce or enlarge** the formula in pharmaceutical preparation. In large manufacturing the official formula must be enlarged, while in the pharmacy or on small products the official formula must be reduced

- **Factor= desired amount/ specified amount**

Examples:

Rx	Codeine phosphate	gr V
	Amaranth solution	ʒ XV
	Alcohol 10 %	f3 ss
	D.W. q.s	f3 I

1. Mitt f3 II

2. Mitt f3 ss

Calculation(1):

$5/15 = 0.3$ g of codeine phosphate

$15/15 = 1$ ml of amaranth

f3 ss = 2 ml f3 = 30 ml f 3 II = 60 ml

factor = $60/30 = 2$

$0.3 \times 2 = 0.6$ g of codeine phosphate

$1 \times 2 = 2$ ml of amaranth

$2 \times 2 = 4$ ml of alcohol

$60 \times 3/4 = 45$ ml

$45 - (4 + 2) = 39$ ml

Procedure(1):

- 1. Weigh 0.6 g of codeine phosphate and put it in a beaker.
- 2. Dissolve the amount of codeine phosphate in 39 ml of D.W.
- 3. Add 2 ml of amaranth and 4 ml of alcohol into the content of the beaker.
- 4. Transfer the content of the beaker into a measuring cylinder and complete the volume to 60 ml by D.W.
- 5. Convert the content of the measuring cylinder into a wide mouth bottle and put a suitable label.

Calculation

- $f_{\text{ss}} = 15 \text{ ml}$ factor = $15 / 30 = 0.5$
- $0.3 \times 0.5 = 0.15 \text{ g}$ of codeine phosphate
- $1 \times 0.5 = 0.5 \text{ ml}$ of amaranth
- $2 \times 0.5 = 1 \text{ ml}$ of alcohol
- $30 \times 0.5 = 15 \text{ ml}$
- $15 \times 3/4 = 11.25 \text{ ml}$
- $11.25 - (0.5 + 1) = 9.75 \text{ ml}$
- Procedure

Follow the same of the above procedure.

Home works

- Rx Calamine 80 g
Zinc oxide 80 g
Glycerin 20 ml
Bentonite magma 250 ml
Calcium hydroxide qs 1000 ml
Mitt gallon

Rx Atropine sulfate gr XX
Camphor water m̄ XV
D.W qs f3 l
Mitt f3 V

A white, cloud-shaped sticker with a small tail at the bottom, containing the text "Thank you!!" in a black, handwritten-style font. The sticker is placed on a brown corkboard background.

Thank
you!!