



Barbiturates

Thiopental Vs. phenobarbital and Dose Calculations

Lab 5

Asst. Lec. Zakariya A. Mahdi , MSc

Department of Pharmacology and Toxicology

University of Al-Mustansiriyah

2020-2021

Overview:

General Anesthesia: Loss of consciousness in addition to loss of sensation


- Analgesia: Loss of sensitivity to pain.
- Sedation: A state of mental calmness, decreased response to environmental stimuli,
- Euthanasia: is the practice of intentionally ending a life to relieve pain and suffering.

Barbiturate


- Class of drugs that act as central nervous system depressants, and can therefore produce a wide spectrum of effects, from mild sedation to total anesthesia
 - Long acting barbiturates, ex. **Phenobarbitone**
 - Short acting barbiturates, ex. **Butobarbitone** and **Pentobarbitone**
 - Ultra short acting barbiturates, ex. **Thiopentane**
- They are also effective as anxiolytics, hypnotics, and anticonvulsants
- Barbiturates also have analgesic effects, however these effects are somewhat weak, preventing barbiturates from being used in surgery in the absence of other analgesics

Mechanism of action

- Enhance binding of GABA with its receptors, prolonged opening of the Chloride channel (influx of Cl), hyperpolarization
- Block the AMPA receptor, a subtype of glutamate receptor, Leading to decrease the activity of excitatory glutamate neurotransmitter



CNS depressant



Barbiturate

- **Advantages:**
 - Rapid anesthetic onset;
 - Provides a prolonged duration of surgical anesthesia.
 - Can be a sedative.
 - Anesthetic agent or euthanasia agent depending on the dose
- **Disadvantages:**
 - Prolonged recovery time.
 - inadequate analgesic properties.
 - extremely expensive.
 - narrow margin of safety.
 - produces respiratory depression at higher dosages.
 - Potent inducer for liver metabolizing enzymes.
 - They have addiction potential, both physical and psychological.

Pharmacokinetics

- High lipid solubility → Cross blood brain barrier, rapid onset
- Redistribution to other tissues → Short duration of action

Experiment protocol

Six mice are injected with Thiopental (three of them By Sc. route and other three by IP route.

Thiopental dosage form :1 gm vial

Thiopental dose in mice:

Thiopental 30-40 mg/kg IP or Sc.

Six mice are injected with Phenobarbital (three of them By Sc. route and other three by IP route.

Phenobarbital dosage form: 200mg/1ml ampoule

Phenobarbital dose in mice:

Pentobarbital 50-90 mg/kg IP

Calculate Your Dose

The dose is the amount of drug taken at any one time

- Weight of drug (e.g. 250 mg)
- Volume of drug solution (e.g. 10 mL, 2 drops)
- The number of dosage forms (e.g. 1 capsule, 1 suppository)
- Other quantity (e.g. 2 puffs)
- **The dosage regimen** is the frequency at which the drug doses are given. Ex. 2.5 mL twice a day, one tablet three times a day...
- Accurate dosing is critical for the proper utilization of all pharmaceuticals
- First you need to know what volume you want to inject into the animal with each treatment being administered, then you need to know how much drug should be in that given volume

Calculate Your Dose

- To calculate the correct dose of drug you need to know
 - The concentration of the drug
 - The weight of the animal
 - The recommended dose rate of the drug for each specific animal model
- **Concentration of the drug**
 - **mg/ml**: Manufacturers usually provide concentrations of their product in milligrams (mg) of drug per (ml) of solvent
 - **%**: 10% solution of Drug A is 10gm/100ml, a 2% solution of Drug A is 2gm/100ml (20mg/ml)
 - **IU/ml**: International Units per ml of, like some of the fat soluble vitamins
 - **powders**: The mg of active drug in the vial. For example, Drug B comes in powdered form with 500mg per vial:
 - If you add 5ml of sterile water for injection to the vial thus providing 5ml of 100mg/ml drug
 - If you add 2.5ml of sterile water for injection, will make 2.5ml of a 200mg/ml solution

Calculate Your Dose

- **Weight of the animal**
 - It is always best to use a scale and get an accurate weight
 - If you cannot weigh the animal prior to injection, you need to be experienced in estimating the weight
- **Dose rate of the drug**
 - Always look up the drug dose for the species you are working with - it often varies

Experimental protocol

- **Parameters**
 - General Activity
 - Characteristics of Breathing
 - **Onset of Sleep (mins)**
 - **Duration of Sleep (mins)**
- Barbiturates are hypnotic drugs
 - Onset of action is the time required to loss the righting reflex
 - Duration of action in mice can be measured by the 'sleeping time' (i.e. the time from the loss of righting reflex to recovery of reflex)

Experimental protocol

- The loss of righting reflex (LORR) assay was used to evaluate sedative/hypnotic effects
- Righting reflex the ability to assume an optimal position when there has been a departure from it
- The onset time of sleep was noted for all animals. After induction of sleep, mice were placed in the inverted position and when sedation was over, the mice came to normal posture and time was noted
- **Record:**
 - .LORR was recorded as the time at which the animal was unable to turn itself (***onset of action***)
 - .The time to regain the righting reflex (***duration of action***)

Results

Group Number	Drug	IP (mouse #1)		Sc (mouse #2)	
		Onset of action (LORR)	Duration of action	Onset of action (LORR)	Duration of action
Subgroup 1	Thiopental				
Subgroup 2	Thiopental				
Subgroup 3	Thiopental				
Subgroup 4	Phenobarbital				
Subgroup 5	Phenobarbital				
Subgroup 6	Phenobarbital				

A decorative graphic in the top-left corner consisting of two parallel lines forming an L-shape. The outer line is dark grey and the inner line is a vibrant blue. The lines are thick and have a slight 3D effect.

Thank you