



LAB NO. 4:

**The tie - line for three
component system**

Done By:

Lecturer Hiba Sabah

Lecturer Zeina Dawood



Ternary systems with one pair of partially miscible liquids:-



- **Water & benzene** are miscible only to a slight extremes so a mixture of the two usually **produces 2 phase system** the lower layer consist of **water saturated with benzene** while the lighter phase (upper) consist of **benzene saturated with water** .



Ternary systems with one pair of partially miscible liquids:-


- On the other hand, alcohol is completely miscible with both benzene & water .it is expected ,therefore, that the addition of sufficient alcohol to 2 phase of water & benzene will **produce single phase system** in which all the three component are miscible.
- **It might helpful to consider alcohol as acting in manner comparable to that of Temperature in binary system of phenol & water .**



- 
- As **heat** used to break the **cohesive** forces between molecules  **miscibility increase** until one phase result ,
 - the addition of alcohol to benzene-water mixture achieves the same but by different means namely **solvent effect** instead of **temperature effect** .
In this case alcohol **serves as intermediate polar solvent** that shifts the **electric equilibrium of the dramatically opposed** highly polar water & non-polar benzene solution to **provide solvation** .



2 component system



3 component system

Alcohol
addition
to:

- water
- benzene

Solubilize the mixture

- One phase





Note:

Adshesive: means force between **Alike** molecules

Cohesive: means force between Like molecules

The tie line with the binodal are *not necessarily parallel* to one another or to the base line as in binary systems, in fact the direction of the tie line are related to the shape of binodal curve, which in turn depends on the **solubility** of the third component (i.e. alcohol) in the other two components .



only when the added component acts *equally* on the other two component to bring them into solution

Will the binodal **perfectly symmetrical** & the tie line run **parallel to the base line** .



Properties of the tie line of three component system :-

1- any system prepared along the tie line both give rise to two phases having a constant composition.

2- the relative amount by weight of the two conjugate phases will depend on the position of the original system along the tie line.



Materials and equipment:-

- 1- H_2O , HAC, CHCl_3 , 1N NaOH solution, phenolphthalein indicator.
- 2- Burette, separatory funnel, conical flask, balance.





Procedure:-

- In a small separatory funnel prepare 50 gm of a mixture having composition giving rise to a two phase system (e.g. 4gm HAC+16 gm CHCl_3 +30gm H_2O).
- **Separate each layer in two conical flasks.**
- Weigh 10 gm for each layer.
- **Titrate each layer with standard 1N NaOH solution using phenolphthalein as indicator. The end point from colorless to pink.**
- Obtain tie line, calculate the percent W/W of HAC in each layer and locate the values on the miscibility curve. **The straight line joining these points should pass through compositions of the two phase system.**



Mixture prepared

(4 ml of HAC) + ($16 \div 1.4 = 11.4$ ml CHCL₃) + (30 ml H₂O)



Calculation:-

- $\text{HAC} + \text{NaOH} \rightarrow \text{NaAC} + \text{H}_2\text{O}$
- 1 M.wt. of HAC = 1 M.wt. of NaOH
- 1 eq.wt of HAC = 1eq.wt of NaOH
- $60 = 1000 \text{ ml } 1\text{N NaOH}$
- $60/1000 = 1 \text{ ml } 1\text{N NaOH}$
- So, each 1 ml of 1N NaOH is equivalent to 0.06 gm, this is the chemical factor (it is the no. of gms of substance which is equivalent to 1 ml of standard solution).
- E.P 1 x 0.06 = gm HAC in 10 gm aqueous layer (upper layer).
- E.P 2 x 0.06 = gm HAC in 10 gm CHCl_3 layer (lower layer).
- Change these values to percent





No. of grams of HAC = E.P(mL of NaOH added) × Ch.F

- Upper layer (between HAC and H₂O)
- wt. of HAC total vol.
- No. of grams 10 g
of HAC
- X 100 g
- X = ? % w/w of HAC
- 100% - X% = ?% w/w
of water

- Lower layer (between HAC and CHCl₃)
- wt. of HAC total vol.
- No. of grams 10 g
of HAC
- X 100 g
- X = ? % w/w of HAC
- 100% - X% = ?% w/w
of CHCl₃



- 
- Upper layer
 - We assume NaOH =
25 mL × 0.06
 - 1.5 gm 10g
 - X 100%
 - X = 15% acetic acid
 - 100% - 15% = 85%
water

- 
- Lower layer
 - We assume NaOH =
8.5 mL × 0.06
 - 0.51 gm 10g
 - X 100%
 - X = 5% acetic acid
 - 100% - 5% = 95%
CHCl₃





X: represent the system

4gm HAC = 8%

16gm CHCl₃ = 32%

30gm H₂O = 60%



Note:-

- For the upper layer represent mostly water with little chloroform.
- This layer represents aqueous layer.
- For the lower layer represent mostly chloroform with little water.
- This layer represents chloroformic layer.

Sp. gr for $\text{CHCl}_3 = 1.4$

Sp. gr for $\text{H}_2\text{O} = 1.009$






Homework lab 4(physical pharmacy)

Prepare 50 gm of a mixture (5gm HAC+20 gm CHCl_3 +25gm H_2O) having composition giving rise to a two phase system, Separate each layer and Weigh 10 gm for each layer then titrate them with standard 1N NaOH solution using phenolphthalein indicator. The end point for the upper layer =28ml and for the lower layer=17 ml,

Q1\calculate the percent W/W of HAC in each layer and locate these values on the miscibility curve to obtain tie line.

Q2\ plot point x (which represent the original system) on the triangle graph paper, Was this point located on the tie line? (Yes or No).





**THANK YOU
FOR YOUR
LISTENING**

