

**OPERATORS : ^**

An operator is a symbol for a mathematical procedure that changes one function into another.

An operator a rule that transforms a given function into another function .

Operators arise because in quantum mechanics you are describing nature with waves ( the wave function ) rather than with discrete particles whose motion can described with the deterministic equations of Newtonian physics.

Operators are more important in classical mechanics and quantum mechanics .

Example 1:

$$D^{\wedge} ( X^2 + 3 e^X ) \text{ if } D^{\wedge} = 3$$

$$\begin{aligned} \text{Solution: } & 3( X^2 + 3 e^X ) \\ & = 3 X^2 + 9 e^X \end{aligned}$$

Example 2 :

$$\begin{aligned} \text{If } D^{\wedge} = d / dx \quad \dots\dots(D^{\wedge} + 3^{\wedge}) ( X^3 - 5) \\ & = (D^{\wedge} X^3 - D^{\wedge} 5) + ( 3^{\wedge} X^3 - 15) \\ & = d/dx X^3 - d/dx 5 + ( 3 X^3 - 15) \\ & = 3X^2 + 3X^3 - 15 \\ & = 3X^3 + 3X^2 - 15 \end{aligned}$$

Types of OPERATORS

1-Commute

$$P^{\wedge}Q^{\wedge}F(x) = Q^{\wedge}P^{\wedge}F(x)$$

### Example 3

$$\text{If } P^\wedge = (d/dx)_{zy}, \quad Q^\wedge = (d/dy)_{zx}$$

sol/

$$\begin{aligned} P^\wedge Q^\wedge F(x) &= (d/dx)_{zy} (d/dy)_{zx} F(x) \\ &= d^2 F(x) / (dx dy)_{xyz} \end{aligned}$$

$$\begin{aligned} Q^\wedge P^\wedge F(x) &= (d/dy)_{zx} (d/dx)_{zy} F(x) \\ &= d^2 F(x) / (dy dx)_{xyz} \end{aligned}$$

2-non commute operators

$$P^\wedge Q^\wedge F(x) \neq Q^\wedge P^\wedge F(x)$$

### Example 4

$$\text{If } P^\wedge = 5+, \quad Q^\wedge = \sqrt{\quad}, \quad F(x) = 4$$

$$P^\wedge Q^\wedge F(x) = 5 + \sqrt{4} = 7$$

$$Q^\wedge P^\wedge F(x) = \sqrt{5+4} = 3$$

### 3- Linear Operators

#### Example4

$$\text{If } F(X) = X^2 + 2x, \quad P^\wedge = d/dx$$

$$\begin{aligned} P^\wedge F(x) &= d/dx (X^2 + 2X) = dX^2/dx + d2X/dx \\ &= 2X + 2 = 2X + 2 \end{aligned}$$

### 4- non-Linear Operators

$$P^\wedge (R+S) \neq P^\wedge R + P^\wedge S$$

#### Example5

$$\text{If } P^\wedge = \sqrt{\quad}, \quad R=5, \quad S=10$$

$$P^\wedge (R+S) = P^\wedge (5+10) = \sqrt{5+10} = \sqrt{15}$$

$$P^\wedge (R+S) = P^\wedge R + P^\wedge S = \sqrt{5} + \sqrt{10}$$

#### Example6

$$\text{If } A^\wedge = d^2/d^2X + 3X d/dX, \quad F(x) = 4X^3$$

$$\begin{aligned} A^\wedge F(x) &= d^2/d^2X + 3X d/dX 4X^3 \\ &= d^2/d^2X 4X^3 + 3X d/dx 4X^3 = 24X + 3X(12X^2) \\ &= 36X^3 + 24X \end{aligned}$$

Example 7

If  $A^{\wedge} = 5$  ,  $F(x) = \sin x$

$$A^{\wedge}F(x) = 5 \sin x$$

Example 8

If  $A^{\wedge} = d/dx$  ,  $F(x) = \cos(X^2+1)$

$$A^{\wedge}F(x) = d/dx \cos(X^2+1)$$

$$= -2X \sin(X^2+1)$$

### Eigen Value Equations

Suppose that effect of operating on some function  $F(x)$  with the operator  $A^{\wedge}$  is simply to multiply  $F(x)$  by a certain constant  $K$  .

We then say that  $F(x)$  is an eigen function of  $A^{\wedge}$  with eigen value  $K$ .

As part of the the definition .we shall require that the eigen function  $F(x)$  is not identically zero.

$$A^{\wedge}F(x) = K F(x)$$

Example 1:

$e^{2x}$  is an eigen function of the operator  $d/dx$  with Eigen value 2

$$d/dx e^{2x} = 2 e^{2x}$$

( operator)(function)=(constant )( same function )

$d/dx =$  operator ,  $e^{2x} =$  eigen function ,  $2 =$  eigen value

Example 2:

If the function  $(\cos ax)$ an eigen function of a)  $d/dx$ , b)  $d^2/d^2x$

Sol/

a)  $d/dx \cos a x = -a \sin ax$ .....non an Eigen function

b)  $d^2/d^2x \cos ax = - a^2 \cos ax$  .....an Eigen function

Example 3:

The function  $X^3$  and the operator  $A^{\wedge} = d^2/ d^2X$  ,  $B^{\wedge} = X^2$  find Eigen function ?

$$\text{Sol/ } A^B X^3 = d^2/d^2x X^2 X^3 = d^2/d^2x X^5 = 20X^3$$

Example 4:

Is wave function  $\sin(z\pi r/a)$  an eigen function for the operator  $(d^2/d^2r) - 4ar^2$ ?

$$\begin{aligned} \text{Sol/ } & (d^2/d^2r - 4ar^2) \sin z\pi r/a \\ & = d^2/d^2r \sin z\pi r/a - 4ar^2 \sin z\pi r/a \\ & = - (z\pi r/a)^2 \sin z\pi r/a - 4 ar^2 \sin z\pi r/a \\ & = - (z\pi r/a)^2 - 4 ar^2) \sin z\pi r/a \dots\dots \text{non Eigen function} \end{aligned}$$

Example 5:

Is wave function  $Y(x) = e^{nx^2}$  an eigen function for the operator  $(d^2/d^2x) - 4n^2x^2$ ?

Sol/

$$\begin{aligned} P^Y(x) &= (d^2/d^2x) - 4n^2x^2) e^{nx^2} \\ &= d^2/dx^2 e^{nx^2} - 4 n^2x^2 e^{nx^2} \\ &= 4 n^2x^2 e^{nx^2} + 2ne^{nx^2} - 4 n^2x^2 e^{nx^2} \\ &= 2ne^{nx^2} \text{ Eigen function} \end{aligned}$$

.....

Q1/ Is wave function  $Q(x) = A e^{-Bx}$  an Eigen function for the operator  $A^{\wedge} = d^2/d^2x + 2/X d/dx + C/X$

Q2/ Is wave function  $f(x) = \sin(-ar)$  an Eigen function for the operator  $A^{\wedge} = d^2/d^2x$

Q3/ Is wave function  $Q(x) = \sin^2 x$  an Eigen function for the operator  $A^{\wedge} = d/dx$

.....