

$$\begin{aligned}
 (x, y) \in A \times (B - C) &\Leftrightarrow x \in A \wedge y \in (B - C) && \text{Def. of } \times \\
 \Leftrightarrow x \in A \wedge (y \in B \wedge y \notin C) &&& \text{Def. of } - \\
 \Leftrightarrow (x \in A \wedge x \in A) \wedge (y \in B \wedge y \notin C) &&& \text{Idempotent Law of } \wedge \\
 \Leftrightarrow (x \in A \wedge y \in B) \wedge (x \in A \wedge y \notin C) &&& \text{Commut. and Assoc. Laws of } \wedge \\
 \Leftrightarrow (x, y) \in (A \times B) \wedge (x, y) \notin (A \times C) &&& \text{Def. of } \times \\
 \Leftrightarrow (x, y) \in (A \times B) - (A \times C) &&& \text{Def. of } -
 \end{aligned}$$

## 3.2 Relations

**Definition 3.2.1.** Any subset “ $R$ ” of  $A \times B$  is called a **relation between  $A$  and  $B$**  and denoted by  $R(A, B)$ . Any subset of  $A \times A$  is called a **relation on  $A$** .

In other words, if  $A$  is a set, any set of ordered pairs with components in  $A$  is a relation on  $A$ . Since a relation  $R$  on  $A$  is a subset of  $A \times A$ , it is an element of the power set of  $A \times A$ ; that is,  $R \subseteq P(A \times A)$ .

If  $R$  is a relation on  $A$  and  $(x, y) \in R$ , then we write  $xRy$ , read as “ $x$  is in  $R$ -relation to  $y$ ”, or simply,  $x$  is in relation to  $y$ , if  $R$  is understood.

**Example 3.2.2.**

(i) Let  $A = \{2, 4, 6, 8\}$ , and define the relation  $R$  on  $A$  by  $(x, y) \in R$  iff  $x$  divides  $y$ . Then,

$$R = \{(2, 2), (2, 4), (2, 6), (2, 8), (4, 4), (4, 8), (6, 6), (8, 8)\}.$$

(ii) Let  $A = \{0, 3, 5, 8\}$ , and define  $R \subseteq A \times A$  by  $xRy$  iff  $x$  and  $y$  have the same remainder when divided 3.

$$R = \{(0, 0), (0, 3), (3, 0), (3, 3), (5, 5), (5, 8), (8, 5), (8, 8)\}.$$

Observe, that  $xRx$  for  $x \in N$  and, whenever  $xRy$  then also  $yRx$ .

(iii) Let  $A = \mathbb{R}$ , and define the relation  $R$  on  $\mathbb{R}$  by  $xRy$  iff  $y = x^2$ . Then  $R$  consists of all points on the parabola  $y = x^2$ .


- (iv) Let  $A = \mathbb{R}$ , and define  $R$  on  $\mathbb{R}$  by  $xRy$  iff  $x \cdot y = 1$ . Then  $R$  consists of all pairs  $(x, \frac{1}{x})$ , where  $x$  is non-zero real number.
- (v) Let  $A = \{1, 2, 3\}$ , and define  $R$  on  $A$  by  $xRy$  iff  $x + y = 7$ . Since the sum of two elements of  $A$  is at most 6, we see that  $xRy$  for no two elements of  $A$ ; hence,  $R = \emptyset$ .

For small sets we can use a pictorial representation of a relation  $R$  on  $A$ : Sketch two copies of  $A$  and, if  $xRy$  then draw an arrow from the  $x$  in the left sketch to the  $y$  in the right sketch.

- (vi) Let  $A = \{a, b, c, d, e\}$ , and consider the relation

$$R = \{(a, a), (a, c), (c, d), (d, b), (d, c)\}.$$

An arrow representation of  $R$  is given in Fig.



- (vii) Let  $A$  be any set. Then the relation  $R = \{(x, x) : x \in A\} = I_A$  on  $A$  is called the **identity relation on  $A$** . Thus, in an identity relation, every element is related to itself only.

**Definition 3.2.3.** Let  $R$  be a relation on  $A$ . Then

- (i)  $\text{Dom}(R) = \{x \in A : \text{There exists some } y \in A \text{ such that } (x, y) \in R\}$  is called the **domain of  $R$** .

- (ii)  $\text{Ran}(R) = \{y \in A : \text{There exists some } x \in A \text{ such that } (x, y) \in R\}$  is called the **range of  $R$** .

Observe that  $\text{Dom}(R)$  and  $\text{Ran}(R)$  are both subsets of  $A$ .

### Example 3.2.4.

(i) Let  $A$  and  $R$  be as in Example 3.2.2.(vi). Then

$$\text{Dom}(R) = \{a, c, d\}, \text{Ran}(R) = \{a, b, c, d\}.$$

(ii) Let  $A = \mathbb{R}$ , and define  $R$  by  $xRy$  iff  $y = x^2$ . Then

$$\text{Dom}(R) = \mathbb{R}, \text{Ran}(R) = \{y \in \mathbb{R} : y \geq 0\}.$$

(iii) Let  $A = \{1, 2, 3, 4, 5, 6\}$ , and define  $R$  by  $xRy$  iff  $x \leq y$  and  $x$  divides  $y$ ;  $R = \{(1, 2), (1, 3), \dots, (1, 6), (2, 4), (2, 6), (3, 6)\}$ , and  $\text{Dom}(R) = \{1, 2, 3\}$ ,  $\text{Ran}(R) = \{2, 3, 4, 5, 6\}$ .

(iv) Let  $A = \mathbb{R}$ , and  $R$  be defined as  $(x, y) \in R$  iff  $x^2 + y^2 = 1$ . Then

$(x, y) \in R$  iff  $(x, y)$  is on the unit circle with centre at the origin. So,

$$\text{Dom}(R) = \text{Ran}(R) = \{z \in \mathbb{R} : -1 \leq z \leq 1\}.$$

### Definition 3.2.5. (Reflexive, Symmetric, antisymmetric and Transitive Relations)

Let  $R$  be a relation on a nonempty set  $A$ .

- (i)  $R$  is **reflexive** if  $(x, x) \in R$  for all  $x \in A$ .
- (ii)  $R$  is **antisymmetric** if for all  $x, y \in A$ ,  $(x, y) \in R$  and  $(y, x) \in R$  implies  $x = y$ .
- (iii)  $R$  is **transitive** if for all  $x, y, z \in A$ ,  $(x, y) \in R$  and  $(y, z) \in R$  implies  $(x, z) \in R$ .
- (iv)  $R$  is **symmetric** if whenever  $(x, y) \in R$  then  $(y, x) \in R$ .

### Definition 3.2.6.

(i)  $R$  is an **equivalence relation** on  $A$ , if  $R$  is reflexive, symmetric, and transitive. The set

$$[x] = \{y \in A : xRy\}$$

is called **equivalence class**. The set of all different equivalence classes  $A/R$  is called the **quotient set**.

(ii)  $R$  is a **partial order** on  $A$  (an **order** on  $A$ , or an **ordering** of  $A$ ), if  $R$  is reflexive, antisymmetric, and transitive. We usually write  $\leq$  for  $R$ ; that is,

$$\boxed{x \leq y \text{ iff } xRy}.$$

- (iii) If  $R$  is a **partial order** on  $A$ , then the element  $a \in A$  is called **least element of  $A$  with respect to  $R$**  if and only if  $aRx$  for all  $x \in A$ .
- (iv) If  $R$  is a **partial order** on  $A$ , then the element  $a \in A$  is called **greatest element of  $A$  with respect to  $R$**  if and only if  $xRa$  for all  $x \in A$ .
- (v) If  $R$  is a **partial order** on  $A$ , then the element  $a \in A$  is called **minimal element of  $A$  with respect to  $R$**  if and only if  $xRa$  then  $a = x$  for all  $x \in A$ .
- (vi) If  $R$  is a **partial order** on  $A$ , then the element  $a \in A$  is called **maximal element of  $A$  with respect to  $R$**  if and only if  $aRx$  then  $a = x$  for all  $x \in A$ .

**Example 3.2.7.**

(i) The relation on the set of integers  $\mathbb{Z}$  defined by

$$(x, y) \in R \text{ if } x - y = 2k, \quad \text{for some } k \in \mathbb{Z}$$

is an equivalence relation, and partitions the set integers into two equivalence classes, i.e., the even and odd integers.

If  $y = 0$ , then  $[x] = \mathbb{Z}_e$ . If  $y = 1$ , then  $[x] = \mathbb{Z}_o$ .  $\mathbb{Z} = \mathbb{Z}_e \cup \mathbb{Z}_o$ ,  $\mathbb{Z}/R = \{\mathbb{Z}_e, \mathbb{Z}_o\}$ .

(ii) The inclusion relation  $\subseteq$  is a partial order on power set  $P(X)$  of a set  $X$ .

(iii) Let  $A = \{3, 6, 7\}$ , and

$$R_1 = \{(x, y) \in A \times A : x \leq y\}, R_2 = \{(x, y) \in A \times A : x \geq y\}$$

$$R_3 = \{(x, y) \in A \times A : y \text{ divisible by } x\}$$

are relations defined on  $A$ .

$$R_1 = \{(3, 3), (3, 6), (3, 7), (6, 6), (6, 7), (7, 7)\},$$

$$R_2 = \{(3, 3), (6, 3), (6, 6), (7, 3), (7, 6), (7, 7)\}.$$

$$R_3 = \{(3, 3), (3, 6), (6, 6), (7, 7)\}.$$

$R_1, R_2$  and  $R_3$  are partial orders on  $A$ .

- (1) The least element of  $A$  with respect to  $R_1$  is .....
- (2) The least element of  $A$  with respect to  $R_2$  is .....
- (3) The greatest element of  $A$  with respect to  $R_1$  is .....
- (4) The greatest element of  $A$  with respect to  $R_2$  is .....
- (5)  $A$  has no least and greatest element with respect to  $R_3$  since, .....
- (6) The maximal element of  $A$  with respect to  $R_3$  is .....
- (7) The minimal element of  $A$  with respect to  $R_3$  is .....

(iv) Let  $X = \{1, 2, 4, 7\}$ ,  $K = \{\{1, 2\}, \{4, 7\}, \{1, 2, 4\}, X\}$  and

$$R_1 = \{(A, B) \in K \times K : A \subseteq B\},$$

$$R_2 = \{(A, B) \in K \times K : A \supseteq B\},$$

are relations defined on  $K$ .

$$R_1 = (\{1,2\}, \{1,2\}), (\{1,2\}, \{1,2,4\}), (\{1,2\}, X),$$

$$(\{4,7\}, \{4,7\}), (\{4,7\}, X),$$

$$(\{1,2,4\}, \{1,2,4\}), (\{1,2,4\}, X),$$

$$(X, X)$$

$$R_2 = (\{1,2\}, \{1,2\}),$$

$$(\{4,7\}, \{4,7\}),$$

$$(\{1,2,4\}, \{1,2\}), (\{1,2,4\}, \{1,2,4\}),$$

$$(X, \{1,2\}), (X, \{4,7\}), (X, \{1,2,4\}), (X, X)$$

$R_1$  and  $R_2$  are partial orders on  $K$ .

- (1)  $K$  has no least element with respect to  $R_1$  since, -----.
- (2) The greatest element of  $K$  with respect to  $R_1$  is -----.
- (3) The least element of  $K$  with respect to  $R_2$  is -----.
- (4)  $K$  has no greatest element with respect to  $R_2$  since, -----.
- (5) The minimal elements of  $K$  with respect to  $R_1$  are -----.
- (6) The maximal element of  $K$  with respect to  $R_1$  is -----.
- (7) The minimal element of  $K$  with respect to  $R_2$  is -----.
- (8) The maximal element of  $K$  with respect to  $R_2$  is -----.

### Remark 3.2.8.

- (i) Every greatest (least) element is maximal (minimal). The converse is not true.
- (ii) The greatest (least) element if exist, it is unique.
- (iii) every finite partially ordered set has maximal (minimal) element.

### Properties of equivalence classes

- (iv) For all  $a \in X, a \in [a]$ .
- (v)  $aRb \Leftrightarrow [a] = [b]$ .
- (vi)  $[a] = [b] \Leftrightarrow (a, b) \in R \Leftrightarrow aRb$ .
- (vii)  $[a] \cap [b] \neq \emptyset \Leftrightarrow [a] = [b]$ .
- (viii)  $[a] \cap [b] = \emptyset \Leftrightarrow [a] \neq [b]$ .
- (ix) For all  $a \in X, [a] \in X/R$  but  $[a] \subseteq X$ .

**Definition 3.2.9.**  $R$  is a **totally order** on  $A$  if  $R$  is a partial order, and  $xRy$  or  $yRx$  for all  $x, y \in A$ ; that is, if any two elements of  $A$  are comparable with respect to  $R$ . Then we call the pair  $(A, \leq)$  a **totally order set** or a **chain**.

### Example 3.2.10.

(i) Let  $A = \{2, 3, 4, 5, 6\}$ , and define  $R$  by the usual  $\leq$  relation on  $\mathbb{N}$ , i.e.  $aRb$  iff  $a \leq b$ . Then  $R$  is a **totally order** on  $A$ .

(ii) Let us define another relation on  $\mathbb{N}$

$$a/b \text{ iff } a \text{ divides } b.$$

To show that  $/$  is a partial order we have to show the three defining properties of a partial order relation:

**Reflexive:** Since every natural number is a divisor of itself, we have  $a/a$  for all  $a \in A$ .

**Antisymmetric:** If  $a$  divides  $b$  then we have either  $a = b$  or  $a < b$  in the usual ordering of  $\mathbb{N}$ ; similarly, if  $b$  divides  $a$ , then  $b = a$  or  $b < a$ . Since  $a < b$  and  $b < a$  is not possible,  $a/b$  and  $b/a$  implies  $a = b$ .

**Transitive:** If  $a$  divides  $b$  and  $b$  divides  $c$  then  $a$  also divides  $c$ . Thus,  $/$  is a partial order on  $N$ .

The relation  $/$  is not totally order since  $(3,4) \notin /$ .

(iii) Let  $A = \{x, y\}$  and define  $\leq$  on the power set  $P(A) = \{\emptyset, \{x\}, \{y\}, A\}$  by

$$s \leq t \text{ iff } s \text{ is a subset of } t.$$

This gives us the following relation:

$$\emptyset \leq \emptyset, \emptyset \leq \{x\}, \emptyset \leq \{y\}, \emptyset \leq \{x, y\} = A, \{x\} \leq \{x\}, \{x\} \leq \{x, y\}, \{y\} \leq \{y\}, \{y\} \leq \{x, y\}, \{x, y\} \leq \{x, y\}.$$

The relation " $\leq$ " is not totally order since  $(\{x\}, \{y\}) \notin \leq$ .

### Exercise 3.2.11.

Let  $A = \{1, 2, \dots, 10\}$  and define the relation  $R$  on  $A$  by  $xRy$  iff  $x$  is a multiple of  $y$ . Show that  $R$  is a partial order on  $A$ .

(Hint:  $R = \{(ny, y) : \text{for some } n \in \mathbb{Z} \text{ and } y \in A\}$ )

**Definition 3.2.12. (Inverse of a Relation)**

Suppose  $R \subseteq A \times B$  is a relation between  $A$  and  $B$  then the inverse relation  $R^{-1} \subseteq B \times A$  is defined as the relation between  $B$  and  $A$  and is given by

$$bR^{-1}a \quad \text{if and only if} \quad aRb.$$

That is,  $R^{-1} = \{(b, a) \in B \times A : (a, b) \in R\}$ .

**Example 3.2.13.** Let  $R$  be the relation between  $\mathbb{Z}$  and  $\mathbb{Z}^+$  defined by  $mRn$  if and only if  $m^2 = n$ .

Then

$$R = \{(m, n) \in \mathbb{Z} \times \mathbb{Z}^+ : m^2 = n\} = \{(m, m^2) \in \mathbb{Z} \times \mathbb{Z}^+\},$$

and

$$R^{-1} = \{(n, m) \in \mathbb{Z}^+ \times \mathbb{Z} : m^2 = n\} = \{(m^2, m) \in \mathbb{Z}^+ \times \mathbb{Z}\}.$$

For example,  $-3 R 9$ ,  $-4 R 16$ ,  $16 R^{-1} 4$ ,  $9 R^{-1} 3$ , etc.

**Remark 3.2.14.** If  $R$  is partial order relation on  $A \neq \emptyset$ , then

- (i)  $R^{-1}$  is also partial order relation on  $A$ .
- (ii)  $(R^{-1})^{-1} = R$ .
- (iii)  $\text{Dom}(R^{-1}) = \text{Ran}(R)$  and  $\text{Ran}(R^{-1}) = \text{Dom}(R)$ .

**Proof. (i)**

(1) **Reflexive.** Let  $x \in A$ .

$$\Rightarrow (x, x) \in R \quad (\text{Reflexivity of } A) \Rightarrow (x, x) \in R^{-1} \quad \text{Def of } R^{-1}$$

(2) **Anti-symmetric.** Let  $(x, y) \in R^{-1}$  and  $(y, x) \in R^{-1}$ . To prove  $x = y$ .

$$\Rightarrow (y, x) \in R \wedge (x, y) \in R \quad \text{Def of } R^{-1}$$

$$\Rightarrow y = x \quad \text{Since } R \text{ is antisymmetric}$$

(3) **Transitive.** Let  $(x, y) \in R^{-1}$  and  $(y, z) \in R^{-1}$ . To prove  $(x, z) \in R^{-1}$ .

$$\Rightarrow (y, x) \in R \wedge (z, y) \in R \quad \text{Def of } R^{-1}$$

$$\Rightarrow (z, y) \in R \wedge (y, x) \in R \quad \text{Commut. Law of } \wedge$$

$$\Rightarrow (z, x) \in R \quad \text{Since } R \text{ is transitive}$$

$$\Rightarrow (x, z) \in R^{-1} \quad \text{Def of } R^{-1}$$

**Definition 3.2.15. (Partitions)**

Let  $A$  be a set and let  $A_1, A_2, \dots, A_n$  be subsets of  $A$  such

- (i)  $A_i \neq \emptyset$  for all  $i$ ,
- (ii)  $A_i \cap A_j = \emptyset$  if  $i \neq j$ ,
- (iii)  $A = \bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \dots \cup A_n$ . Then the sets  $A_i$  partition the set  $A$  and these sets are called the **classes of the partition**.

**Remark 3.2.16.** An equivalence relation on  $X$  leads to a partition of  $X$ , and **vice versa** for every partition of  $X$  there is a corresponding equivalence relation.

**Proof:**

(a) Let  $R$  be an equivalence relation on  $X$ .

1-  $\forall a \in X, a \in [a]$  Def. of equ. Class

2-  $\exists [b] \in X/R$  such that  $[b] = [a]$  Since  $X/R$  contains all diff. classes

3-  $X = \bigcup_{a \in X} \{a\} \subseteq \bigcup_{a \in X} [a] \subseteq \bigcup_{a \in [b]} [b] \subseteq X \Rightarrow X = \bigcup_{[b] \in X/R} [b]$ .

4-  $[b] \cap [a] = \emptyset$ , for all  $[b], [a] \in X/R$  Def. of  $X/R$

5-  $R$  is partition of  $X$  Inf.(3),(4)

(b) Let (i)  $A_i \neq \emptyset$  for all  $i, A_i \subseteq X$

(ii)  $A_i \cap A_j = \emptyset$  if  $i \neq j$ ,

(iii)  $X = \bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \dots \cup A_n$ .

Define  $R$ (relation) on  $X$  by  $aRb \Leftrightarrow$  if  $\exists A_i$  such that  $a, b \in A_i$ .

This relation is an equivalence relation on  $X$ .

**Definition 3.2.17. (The Composition of Two Relations)**

The composition of two relations  $R_1(A, B)$  and  $R_2(B, C)$  is given by  $R_2 \circ R_1$  where  $(a, c) \in R_2 \circ R_1$  if and only if there exists  $b \in B$  such that  $(a, b) \in R_1$  and  $(b, c) \in R_2$ . That is,

$$R_2 \circ R_1 = \{(a, c) \in A \times C \mid \exists b \in B \text{ such that } (a, b) \in R_1 \text{ and } (b, c) \in R_2\}$$



**Remark 3.2.18.** Let  $R_1(A, B)$ ,  $R_2(B, C)$  and  $R_3(C, D)$  are relations. Then,

(i)  $(R_3 \circ R_2) \circ R_1 = R_3 \circ (R_2 \circ R_1)$ .

(ii)  $(R_2 \circ R_1)^{-1} = R_1^{-1} \circ R_2^{-1}$ .

(iii) Let  $R^{-1} = \{(b, a) | (a, b) \in R\} \subseteq B \times A$ . Then

$$(a, b) \in R \circ R^{-1} \Leftrightarrow (b, a) \in R \circ R^{-1}.$$

**Proof. Exercise.**

**Example 3.2.19.**

Let sets  $A = \{a, b, c\}$ ,  $B = \{d, e, f\}$ ,  $C = \{g, h, i\}$  and relations


$$R(A, B) = \{(a, d), (a, f), (b, d), (c, e)\}$$


and

$$S(B, C) = \{(d, g), (d, i), (e, g), (e, h)\}.$$

Then we graph these relations and show how to determine the composition pictorially  $S \circ R$  is determined by choosing  $x \in A$  and  $y \in C$  and checking if there is a route from  $x$  to  $y$  in the graph. If so, we join  $x$  to  $y$  in  $S \circ R$ .

$$S \circ R = \{(a, g), (a, i), (b, g), (b, i), (c, g), (c, h)\}.$$





For example, if we consider  $a$  and  $g$  we see that there is a path from  $a$  to  $d$  and from  $d$  to  $g$  and therefore  $(a, g)$  is in the composition of  $S$  and  $R$ .

### Definition 3.2.19. Union and Intersection of Relations

(i) The union of two relations  $R_1(A, B)$  and  $R_2(A, B)$  is subset of  $A \times B$  and defined as

$$(a, b) \in R_1 \cup R_2 \text{ if and only if } (a, b) \in R_1 \text{ or } (a, b) \in R_2.$$

(ii) The intersection of two relations  $R_1(A, B)$  and  $R_2(A, B)$  is subset of  $A \times B$  and defined as

$$(a, b) \in R_1 \cap R_2 \text{ if and only if } (a, b) \in R_1 \text{ and } (a, b) \in R_2.$$

### Remark 3.2.20.

(i) The relation  $R_1$  is a subset of  $R_2$  ( $R_1 \subseteq R_2$ ) if whenever  $(a, b) \in R_1$  then  $(a, b) \in R_2$ .

(ii) The intersection of two equivalence relations  $R_2, R_1$  on a set  $X$  is also equivalence relation on  $X$ .

(iii) In general, the union of two equivalence relations  $R_1, R_2$  on a set  $X$  need not to be an equivalence relation on  $X$ .

### Proof. Exercise.

**Example 3.2.21.** Let  $X = \{a, b, c\}$ . Define two relations on  $X$  as follows:

$$R_1(X, X) = \{(a, a), (b, b), (c, c), (a, b), (b, a)\},$$

$$R_2(X, X) = \{(a, a), (b, b), (c, c), (a, c), (c, a)\}.$$

Let  $R = R_1 \cup R_2$ . Here,  $R$  is not an equivalence relation on  $X$  since it is not transitive relation, because  $(b, a)$  and  $(a, c) \in R$  but  $(b, c) \notin R$ .

Dr. Bassam and Dr. Emad