

# LAB. METEOROLOGICAL STATISTICS ..... FOURTH STAGE

(First Semester)

Department of Atmospheric Sciences

2021 – 2022

**Lecturers: L. Ruaa mazin , A.L. Yasamin qusay , A.L. Zahraa araf , A.L. Luma Mahdi , A.L. Salwa salman**

**Preparing by: L. Ruaa mazin , A.L. Zahraa araf , A.L. Luma mahdi**

## Measures of Dispersion and Variability

measure of variability is a summary statistic that represents the amount of dispersion in a dataset. How spread out are the values? A low dispersion indicates that the data points tend to be clustered tightly around the center. High dispersion signifies that they tend to fall further away.

### ***A- Range:***

Let's start with the range because it is the most straightforward measure of variability to calculate and the simplest to understand. The range of a dataset is the difference between the largest and smallest values in that dataset.

***For example,*** \\ calculate the range of the dataset: -

60,65.55,50,45

$$\text{Range} = 65 - 45 = 20$$

***For example*** \\ calculate the range from the dataset: -

100,65,55,50,45

$$\text{Range} = \max - \min = 100 - 45 = 55$$

***For example,*** // The numbers are 7,4,9,7,3,12 find the range?

**Solute** \\

$$\text{Range} = 12 - 3 = 9$$

H.W\ Calculate the Rang in the two datasets below

<u>datasets1</u>	<u>datasets2</u>
20	11
21	16
22	19
25	23
26	25
29	32
33	39
34	46
38	52

***B-Mean deviation(MD):***

The mean deviation is defined as a statistical measure that is used to calculate the average deviation from the mean value of the given data set. The mean deviation of the data values can be easily calculated using the below procedure:

Step 1: Find the mean value for the given data values

Step 2: Now, subtract the mean value from each of the data values given (Note: Ignore the minus symbol)

Step 3: Now, find the mean of those values obtained in step 2.

**1- The mean deviation of the unclassified data**

The mean deviation or average deviation of a sat N numbers  $X_1, X_2, X_3, \dots, X_N$

Is MD and defined by:

$$\text{Mean deviation (MD)} = \frac{\sum |x - \bar{x}|}{N}$$

$\Sigma$  represents the addition of values

X represents each value in the data set

$\bar{x}$  represents the mean of the data set (arithmetic mean)

N represents the number of data values

|| represents the absolute value, which ignores the “-” symbol

**For example** // find the mean of the set : 2,3,6,8,11

$$\bar{x} = \frac{\Sigma x_i}{N} = \frac{2+3+6+8+11}{5} = \frac{30}{5} = 6$$

$$MD = \frac{|2-6|+|3-6|+|6-6|+|8-6|+|11-6|}{5}$$

$$MD = \frac{|-4|+|-3|+|0|+|2|+|5|}{5} = \frac{4+3+0+2+5}{5} = \frac{14}{5} = 2.8$$

**H.W\** Calculate the mean deviation from the following data:

6,7,10,12,13,4,8,12

**2- the Mean deviation for the classified data**

The mean deviation can be given by: -

$$MD = \frac{\Sigma f |x - \bar{x}|}{\Sigma f}$$

Where (f) is frequency.

(x) is midpoint of classes.

For example, // find the mean deviation of the following data

<u>Class</u>	<u>Frequency(fi)</u>	<u>Midpoint(xi)</u>	<u>fixi</u>
10-20	2	$10+20/2=15$	$2*15=30$
20-30	3	25	75
30-40	8	35	280
40-50	14	45	630
50-60	8	55	440
60-70	3	65	194
70-80	2	75	150
	$\sum fi=40$		$\sum fixi=1800$

We find the arithmetic mean

$$\bar{x} = \frac{\sum f_i x_i}{\sum f_i} = \frac{1800}{40} = 45$$

<u><math> x_i - \bar{x} </math></u>	<u>Abs</u>	<u>f</u>	<u><math>f *  x_i - \bar{x} </math></u>
$ 15 - 45 $	$ -30 $	2	60
$ 25 - 45 $	$ -20 $	3	60
$ 35 - 45 $	$ -10 $	8	80
$ 45 - 45 $	$ 0 $	14	0
$ 55 - 45 $	$ 10 $	8	80
$ 65 - 45 $	$ 20 $	3	60
$ 75 - 45 $	$ 30 $	2	60

$$\sum f * |x_i - \bar{x}| = 400$$

$$MD = \frac{\sum f |x - \bar{x}|}{\sum f} = \frac{400}{40} = 10$$

H.W\ Calculate the mean deviation from the following data:

<i><u>classes</u></i>	<i><u>f</u></i>
2-4	2
4-6	3
6-8	6
8-10	2
10-12	1