



(PG)

Physical Chemistry_Chpt_One_Properties of Gases

50/100 Fifty only
I.R. Rushdi

Name of a student Teba Hussein Alwi Signature _____ No. 30

University of Mustansiriyah

1st Semester-2021

Department of Chemistry

1st Exam-paper E

Q1: Circle the right answer for all of the following:

(50 points)

1: If a gas has polar particles then the difference between the volume of this gas is:

Answer: a) $V_{Real} > V_{Perfect}$ b) $V_{Real} < V_{Perfect}$ c) $V_{Real} = V_{Perfect}$ d) $V_{Real} \neq V_{Perfect}$

2: A gas occupies 60×10^3 mL at 150°C and 760 mmHg pressure. What would be its volume at STP?

Answer: a) 38.7 mL b) 38.7 dm^3 c) 38.7 L^{-1} d) 38.7 dm^{-3}

3: Calculate the weight of H_2O gas (18 g.mol^{-1}) in a 5 L cylinder at 10×10^2 kPa and 373 K.

Answer: a) 29.40 g mol^{-1} b) 29.40 g c) 29.40 mol d) 29.40 kg

4: Calculate the density of H_2O placed in a 22400 mL cylinder at 10^5 Pa and 0°C .

Answer: a) 0.804 kg L^{-1} b) 0.804 g L^{-1} c) 0.804 g d) 0.804 L^{-1}

5: According to Graham's law the heaviest gas is?

Answer: a) H_2O b) CH_4 c) NH_3 d) Cl_2

6: A tank contains a certain amount of gas at 10^5 Pa. The gas is transferred to another tank 40 dm^3 with pressure of 200×10^3 Pa. What should be its volume?

Answer: a) 80 L b) 80 Pa L c) 80 Pa dm^3 d) 80 L^{-1}

7: According to Boyle's law the pressure of a gas is inversly proportional with?

Answer: a) p b) T c) R d) V e) n

8: The difference between real and ideal gas, that the real gas interested in?

Answer: a) V & p b) V & T c) p & n d) T & p

9: It can follow the direct proportional between temperature and pressure through the law of

Answer: a) Van der Waal b) Graham c) Charles d) Gay-Lussac

10: The behaviour of real gas is ideal when the value of Z is equal to

Answer: a) $V_m < V_m^0$ b) $V_m > V_m^0$ c) $V_m = V_m^0$ d) $V_m \neq V_m^0$

Q2: The following data have been observed for 800 mg of nitrogen gas at 273 K. Calculate the best value of the molar mass of N_2 .

$p/10^5 \text{ Pa}$	0.750	0.500	0.200
V/dm^3	3.0	4.5	7.0

(25 points)

Q3: A perfect gas undergoes isothermal compression, which reduces its volume by 1.80 dm^3 . The p_f and V_f of the gas are 2×10^2 kPa and 2.14 dm^3 , respectively. Calculate the p_{original} of the gas in (i) bar, (ii) torr. (25 points)

Thur_11/11/2021

Best wishes

Dr Abduljabbar I. R. Rushdi

bar =
torr =

92

~~PV = nRT~~

~~PV = $\frac{m}{M} RT$~~

~~M = $\frac{mPV}{RT}$~~

~~M = $\frac{0.75 \times 3}{0.82 \times 273}$~~

~~M = $\frac{2.25}{22.386} = ?$~~

Q2 $\frac{5}{25}$

93

~~P₁V₁ = P₂V₂~~

its not ~~V~~

~~P₁ × 1.80 dm³ = 2 × 10² kPa × 2.14 dm³~~

~~P₁ × 1.80 dm³ = 4.28 × 10² kPa · dm³~~

~~P₁ = $\frac{4.28 \times 10^2 \text{ kPa} \cdot \text{dm}^3}{1.80 \text{ dm}^3}$~~

Q3 $\frac{10}{25}$

~~P = 2.377 × 10² kPa~~

is not P, its bar
↓
bar

~~1 bar = 101.325 kPa ⇒ $\frac{2.377 \times 10^2}{101.325} = 2.343 \text{ bar} = 0.023 \times 10^2 \text{ bar}$~~

~~Q4~~

~~2.33 × 10² kPa × 760 torr = 1806.52^{10^2} torr~~