



0-5 Mid Quiz

Physical_Chemistry_2nd_YUGS_EV_ST

Name of a student

Signature

No. 5

Mustansiriyah University
Department of Chemistry

2nd SEM-2025_Bologna_Process
Mid_Exam_Class_A_Paper_C

Q1/MCO test (Answer the following)

Q. 50

(Marks 50 %)

1: Depression of freezing point of a solution means increasing in?

- Answer: a) T b) H c) μ d) S

2: If you apply the reduced phase rule to condensed systems, then the expected value of pressure is -----?

- Answer: a) zero b) 1 c) 2 d) 3

3: The reduced phase rule can be applied when the number of components is -----?

- Answer: a) zero b) 1 c) 2 d) 3

4: Which One of the following formulas represents the right equation of negative deviation from Raoult's law?

- Answer: a) $P_A^* \neq \chi_A P_A$ b) $P_A = \chi_A P_A^*$ c) $P_A > \chi_A P_A^*$ d) $P_A < \chi_A P_A^*$

5: Addition of a non-volatile solute to the pure solvent causes a change in?

- Answer: a) $\Delta_{mix}H$ b) $\Delta_{mix}S$ c) $\Delta_{mix}V$ d) all of these

6: The difference between pure and impure solvent is?

- Answer: a) $\mu^* = \mu$ b) $\mu^* > \mu$ c) $\mu^* < \mu$ d) $\mu^* \neq \mu$

7: The relationship between ΔT_f and χ_B is?

- Answer: a) direct b) inverse c) disordered d) none of these

8: With the two-component system (A & B), one part of the solid phase consists of?

- Answer: a) A + B b) A + solution c) B + solution d) A + eutectic

9: If you add a solute to a solvent, then there is a decrease in the ----- of the solution.

- Answer: a) S b) H c) T d) μ

10: Dalton's law is used to calculate the partial pressure of ----- phase?

- Answer: a) liquid b) gas c) solid d) plasma

Q2 The Π of a solution containing 4.0 g of an unknown substance per 0.5 dm³ of solution is 10³ torr at

34.0 °C. Find the molar mass of this unknown.

(Marks 25%)

Q3 Using the diagram below and the appropriate phase rule, fill in all the blanks and determine the composition of the all-eutectic mixture, all equilibria, all reversible and irreversible processes, and the name of the regions located to the right and left of points C, E & AB? (Marks 25%)



Name of a student _____ Signature _____ No. --

$$\textcircled{1} \quad T = 273 + 34.0^\circ\text{C} = 307 \text{ K} \quad ? \equiv \text{Units}$$

$$\text{Th} = RT[B] \Rightarrow \textcircled{8} = 0.082 \times 307 [\text{B}]$$


$$\textcircled{9} \quad 0.082 = 25.17 [\text{B}] \Rightarrow \text{B} = \frac{0.082}{25.17}$$

$$[\text{B}] = 0.317 \text{ M}$$

$$\text{so fmolar} = M \times V^{\text{wt}} \Rightarrow 0.317 \times 0.5 = 0.158$$

$$\text{molar mass} = \frac{\text{wt}}{\text{molar mass}} = \frac{4}{0.158} = 26.29 \text{ g/mol}$$

Two component system ()



النقطة A هي النقطة المطلوبة في المحلول A-B
لذلك تكون النتيجة A-B