



# Mid Quiz

Physical\_Chemistry\_2<sup>nd</sup>\_YUGS\_EV\_ST

Name of a student ميسى جعفر Signature Patra No. 33

Mustansiriyah University  
Department of Chemistry

100  
100  
Ten Only  
18-03-25  
Date  
2<sup>nd</sup> SEM 2025 Bologna Process  
Mid Exam Class A Paper D

## Q1/ MCQ test (Answer the following)

(Marks 50 %)

1: The Gibbs phase rule is interested in two variants?

Answer: a) p & T      b) F & T      c) p & conc.

d) T & conc.

2: What do you expect if you add NaCl to H<sub>2</sub>O, an increase in the?

Answer: a) LP      b) VP      c) FP

d) BP

3: The three phases of CO<sub>2</sub> in the phase diagram meets?

Answer: a) at 1 atm      b) over 1 atm      c) below 1 atm

d) at any pressure

4: The phase of super cooling is -----?

Answer: a) gas      b) liquid      c) solid

d) plasma

5: How many phases are there when the number of variants is two and the number of components is one?

Answer: a) zero      b) 1      c) 2      d) 3

6: The Clapeyron equation can be applied when there is an equilibrium between one of the following?

Answer: a) melt. & freez.      b) sub. & depo.      c) vap. & cond.

d) all of these

7: The relationship between VP and m is -----.

Answer: a) direct      b) inverse      c) disordered      d) none of these

8: If you add a ----- to a solvent, then there is a change in the colligative properties of the solvent.

Answer: a) non-volatile solute      b) volatile solute      c) pure solute

d) pure solvent

9: Osmotic process is used to push the solvent to the -----?

Answer: a) solute      b) impure solute      c) mixture      d) pure solvent

10- One of the most important benefits of measuring molar mass of the solute is to study the change in -----.

Answer: a) m      b) Π      c) V      d) p

Q2/ 0.5 mol of a non-P-solute was added to 12.0 mol of P-solvent, VP\* is 12.0 kPa at 295 K. What is the VP at

295 K? Determine the deviation of this solution from Raoult's law where VP<sub>ideal</sub> = 10 kPa. (Marks 25%)

Q3/ Plot the phase diagram of the system (A & B) assumed that (A & B) do not react with each other. A

freezes at (-6 °C) and B freezes at (8 °C), and that an eutectic mixture is formed when the ratio is 60 wt

% of A and that the eutectic melts at (-9 °C), then label all the parts (p & F) of the diagram? (Marks 25%)

~~$P_A = X_A P_A^*$~~   
 ~~$= 10 - 5 \text{ m}_0 L (12 \text{ kPa})$~~

$P_A = 6$

$V_{\text{ideal}} = 70 \text{ kPa}$   
 $P_A = 6$   
 فرق درجة حرارة يكون أكثر من  $P_B$  غير المفترض

