



Physical_Chemistry_2nd_YUGS_EV_ST

25/100 Twenty five



Name of a student _____ Signature _____ No. _____

Mustansiriyah University
Department of Chemistry

1st SEM-2025_Bologna_Process
Mid_Exam_Class_A_Paper_A

Q1: Circle the right answer for all of the following

(50 Marks)

1: liquefaction of the gas means which of the following?

- (a) $Z = 1$
- (b) $Z > 1$
- (c) $Z < 1$
- (d) $Z \neq 1$

2: In the van der Waals equation, what is the correct formula for the volume of the gas?

- (a) n_i/n_T
- (b) V
- (c) V/m
- (d) V/n

3: If a gas has polar particles, then the difference between the volume of this gas is:

- (a) $V_{Real} > V_{Perfect}$
- (b) $V_{Real} < V_{Perfect}$
- (c) $V_{Real} = V_{Perfect}$
- (d) $V_{Real} \neq V_{Perfect}$

4: It can classify the type of reaction within adiabatic process as:

- (a) reversible
- (b) isobaric
- (c) isothermal
- (d) free expansion

5: If it is required to measure the work done in an isochoric process, the value of work will be:

- (a) zero
- (b) one
- (c) two
- (d) three

6: In a completely insulated system, the work done is in contact with which of the following?

- (a) gas
- (b) system
- (c) surrounding
- (d) pressure

7: When the internal pressure of the system is equal to atmospheric pressure, the actual value will be:

- (a) zero
- (b) one
- (c) two
- (d) three

8: During an isothermal reversible process, the change in temperature of the system is?

- (a) variable
- (b) equal to zero
- (c) 25°C
- (d) constant

9: Heat capacity is extensive property while molar heat capacity is ----- property:

- (a) proportional
- (b) intensive
- (c) extensive
- (d) direct

10: When $\Delta H = \text{zero}$, the process is:

- (a) isobaric
- (b) isochoric
- (c) isothermal
- (d) adiabatic

Q2: Using van der Waals equation, calculate the temperature of 5.0 mol of an unknown gas in a 5.0 L container at 80 bar. Compare this temperature with the value obtained from the ideal gas equation.

$a = 0.0341 \text{ L}^2 \text{ atm mol}^{-2}$; $b = 0.0237 \text{ L mol}^{-1}$.

(25 Marks)

Q3: 1100 J of heat energy was applied to (50 g, 27 g mol^{-1}) of aluminum metal. The temperature increased from 25°C to 45°C . Calculate $C_{p,m}$ of aluminum.

(25 Marks)

Q2

~~PV = nRT~~ → ~~PV = $\frac{w}{M \cdot w_t} RT$~~

~~(0.5)(0.5) = $\frac{0.0341}{0.0237} (0.082) 80$~~

Zero
Q2 25

Q3

~~PV = nRT~~ → ~~PV = $\frac{w}{M \cdot w_t} RT$~~

~~(1100) 50 - 27 =~~ Wrong eq!

Zero
Q3 25