



Handwritten marks: 1/5

F10

Physical Chemistry 2<sup>nd</sup> YUGS\_EV\_ST

Handwritten marks: 20/20, 10/10, 10/10



Name of a student: \_\_\_\_\_ Signature: \_\_\_\_\_ No. B2  
Mustansiriyah University  
Department of Chemistry

2<sup>nd</sup> SEM-2026\_Bologna\_Process  
Mid\_Exam\_Class\_A\_Paper\_B

Q1/ MCO test (Answer the following)

Marks 50 %

1: Which two variables does the Gibbs phase rule consider as independent??

Answer: a) p & T b) F & T c) p & conc. d) T & conc.

2: If NaCl is added to ice, which property of the solution decreases?

Answer: a) LP b) VP c) FP d) BP

3: At what pressure do the three phases of CO<sub>2</sub> coexist in the phase diagram?

Answer: a) at 1 atm b) over 1 atm c) below 1 atm d) at any pressure

4: Which phase corresponds to a supercooled substance?

Answer: a) gas b) liquid c) solid d) plasma

5: How many phases are present when a one-component system has two degrees of freedom?

Answer: a) zero b) 1 c) 2 d) 3

6: The Clausius equation can be applied to which of the following phase equilibria?

Answer: a) melt. & freez. b) freez. & melt. c) vap. & cond. d) all of these

7: What is the relationship between the VP of a solution and the solute molality?

Answer: a) direct b) inverse c) disordered d) none of these

8: Which type of solute, when added to a solvent, alters its colligative properties?

Answer: a) non-volatile solute b) volatile solute c) pure solute d) pure solvent

9: In osmosis, the solvent moves toward which component?

Answer: a) solute b) impure solute c) mixture d) pure solvent

10- One of the most important applications of measuring molar mass of the solute is to study the change in ---.

Answer: a) m b)  $\Pi$  c) V d) p

Q2/ 0.5 mol of a non-P-solute was added to 12.0 mol of P-solvent, VP\* is 12.0 kPa at 295 K. What is the VP at

295 K? Determine the deviation of this solution from Raoult's law where VP<sub>ideal</sub> = 10 kPa. (Marks 25%)

Q3/ Plot the phase diagram of the system ( $\alpha$  and  $\beta$ ) assumed that ( $\alpha$  and  $\beta$ ) do not react with each other.  $\alpha$

freezes at (-7 °C) and  $\beta$  freezes at (10 °C), and that a eutectic mixture is formed when the ratio is 30 wt % of  $\beta$  and that the eutectic melts at (-10 °C), then label all the parts (p & F) of the diagram using the

appropriate phase rule?

(Marks 25%)

Q2/

$$0.5 + 12 = 12.5 \text{ mol}$$

بدرجه حراره 295K

Where is the eq?

0.5 mol من غير مقياس

باصفايته 12 mol

بدرجه حراره 295K

الفترت بيننا وبين مقياس P بالاولت

وضيحه VP را اولت = 10kPa

$$12.5 - 10 = 2.5 \text{ mol}$$

بدرجه حراره 295K

سيكون 12.5 اجل من مقياس اولت

Wrong info!


Zero  
Q2/25

Q3/

~~Water~~

~~ferol~~  
~~سول~~

Being point 4  
Tempr



~~Solid, Liquid~~

~~Liquid, gas~~

~~P=1~~

~~C=2~~

~~P=1~~

~~C=2~~

0%  
100%

100%  
0%